

# Updates in transition cow health and nutrition

Thomas R. Overton,<sup>1</sup> MS, PhD; Sabine Mann,<sup>2</sup> DVM, PhD; Brittany M. Leno<sup>1</sup>; Daryl V. Nydam,<sup>2</sup> DVM, PhD

<sup>1</sup>Department of Animal Science, Cornell University, Ithaca, NY 14853

<sup>2</sup>Department of Population Medicine and Diagnostic Sciences, Cornell University, Ithaca, NY 14853

## Abstract

Transition cow nutritional management strategies have evolved significantly over the past 10 years and continue to evolve. Management of subclinical hypocalcemia is important to support high milk yield, excellent reproduction, and decrease subclinical disease. Feeding anionic diets before calving, perhaps using more aggressive strategies, improves blood calcium status post-calving and increases post-calving dry matter intakes and milk yield. Controlling energy intake before calving improves postpartum energy balance and decreases subclinical ketosis; feeding adequate metabolizable protein before calving helps maintain high milk yield after calving. Excellent feeding management (managing chop length of bulky forage and moisture of the (total mixed ration) TMR) of transition rations is critical for on-farm success; many dairy farms have opportunities for improved feeding management. There may be interactions of starch levels pre- and post-calving; cows fed low starch pre-calving rations should be fed lower starch diets after calving, whereas cows fed moderate starch rations pre-calving can likely be fed starch levels more typical of high starch cow rations after calving. Furthermore, there may be opportunities to combine higher fermentability with higher bulk in fresh cow rations to improve feed intakes after calving, as well as to more critically consider metabolizable protein and amino acid formulation of the fresh diet.

**Key words:** dairy, transition, feeding, health

## Résumé

Stratégies de gestion de la nutrition des vaches en transition ont évolué considérablement au cours des 10 dernières années et continuera d'évoluer. Gestion de l'hypocalcémie subclinique est important de soutenir la production de lait, une excellente reproduction, et la diminution des maladies subcliniques. Les régimes alimentaires anioniques d'alimentation avant le vêlage, peut-être en utilisant des stratégies plus agressives, améliore l'état de calcium de sang après la mise bas et de post-mise bas augmente l'apport de matière sèche et la production de lait. Contrôler l'apport en énergie avant le vêlage améliore le bilan énergétique postpartum et la cétose subclinique diminue ; les protéines métabolisable adéquate avant le vêlage permet de maintenir un haut niveau de production laitière après le vêlage. Excellente gestion de l'alimentation (la gestion de la durée de broyage du fourrage encombrant et de l'humidité de la ration totale

mélangée (RTM)) de rations de transition est cruciale pour le succès à la ferme ; de nombreuses fermes laitières ont des possibilités d'amélioration de l'alimentation de la gestion. Il peut y avoir des interactions entre les niveaux d'amidon pré- et post-mise bas ; vaches nourries bas amidon pré-mise bas devraient être nourries de rations alimentation amidon inférieure après le vêlage, tandis que vaches nourries de rations d'amidon modérée vêlage pré-peuvent probablement être nourries plus typiques des niveaux d'amidon l'amidon des rations de vache après le vêlage. En outre, il peut être possible de combiner des fermentation avec des rations de vache frais en vrac pour améliorer les prises d'alimentation après le vêlage, ainsi qu'à examiner de façon plus critique et d'acides aminés protéines métabolisable de formulation de l'alimentation.

## Introduction

Nutritional management strategies for transition cows have evolved significantly over the past 10 years in the dairy industry, and ongoing research and experience continue to refine our recommendations for nutritional management of dairy cows during both the prepartum and postpartum periods. Our high-performing herds combine high milk production with modest loss of body condition score (no more than 0.50 or so units during early lactation), low incidence of metabolic and immune function-related diseases, excellent reproductive performance during early lactation, and have calves born alive and ready to thrive. In this paper, we will outline several areas that have received significant research attention during the past several years, along with some that are currently very active areas of research that likely will lead to continued evolution of nutritional recommendations for transition cows.

## Feeding Strategies for the Dry Cow

*Management of hypocalcemia – an old topic made new again*

Clinical milk fever is a thing of the past on many dairies. Research over the past 5 years has shifted the hypocalcemia focus to include management of not only clinical cases of milk fever, but also cows that experience subclinical drops in blood calcium postpartum. Even in herds with very low milk fever incidence, subclinical hypocalcemia (SCH) after calving can affect 50% or more of the herd, predisposing cows to infectious and metabolic disease and reducing their productive and reproductive potential.<sup>4,18,21</sup> As these associations continue to be researched, the need for strategies to

reduce SCH incidence is becoming more evident. Reducing the dietary cation anion difference (DCAD;  $\text{Na} + \text{K} - \text{Cl} - \text{S} = \text{mEq}/100 \text{ g DM}$ ) of the prepartum ration is a tried and true method for decreasing rates of clinical milk fever.<sup>2,9</sup> Strategies for implementing this approach can range from minimizing the dietary potassium (aiming for a low but still positive DCAD) to varying inclusion rates of anion supplements to reach a negative DCAD.

Recent work by our group at Cornell University aimed to determine if benefits in calcium status and production parameters increased when anion inclusion rate was incrementally increased (and therefore, DCAD decreased) in a low-potassium prepartum ration.<sup>22,23</sup> Three experimental groups included a low-potassium control ration (+18.3 mEq/100 g DM), partial anion supplementation (+5.9 mEq/100 g DM), and full anion supplementation (-7.4 mEq/100 g DM). Diets were managed to maintain urine pH of the full anion supplemented group between the target of 5.5 to 6.0. Ultimately, as prepartum DCAD was decreased in this trial, average postpartum plasma calcium was increased, indicating that the greatest benefit in calcium status postpartum was seen in cows fed the lowest DCAD. Interestingly, an effect of parity was seen such that older cows (3rd+ lactation) benefited the most when fed the lowest DCAD.<sup>22</sup> Increases in postpartum dry matter intake and milk yield were seen for cows fed decreasing DCAD. Cows fed the lowest DCAD ration prepartum produced over 7 lb (3.2 kg)/day more milk in the first 21 days compared to cows fed the low-potassium control ration.<sup>23</sup> This study indicates that implementing a more aggressive DCAD prepartum can yield the greatest benefits postpartum when compared to a low-potassium control approach.

Measuring urine pH is an essential component of monitoring prepartum DCAD, and can also provide valuable information about feeding management.<sup>5,11</sup> Urine pH should be measured in midstream urine samples from approximately 12 to 15 cows weekly. It is important that the time relative to feeding is consistent from week to week, since urine pH response may fluctuate throughout the day. Large variation from cow to cow within a week may indicate undesirable consumption of the ration, whether that be a result of overcrowding, social factors, or sorting due to poor diet mixing. Variation in average urine pH from week to week can indicate inconsistency in ration mixing or changes in feed ingredient composition. This information can be used to improve feeding and management strategies to increase transition cow success.

#### *Dry period plane of energy and effects on health, production, and reproduction*

Since the early 2000s we have largely abandoned the historically proposed “steam up” approach to dry cow feeding. With increasing evidence from research conducted at the University of Illinois, a controlled energy strategy to feeding dry cows was proposed.<sup>8</sup> Lower postpartum concentrations of non-esterified fatty acids (NEFA) and ketone bodies (e.g.

$\beta$ -hydroxybutyrate or BHB) were observed with these controlled energy diets and the incidence of metabolic disease was decreased.<sup>6,10</sup> However, detrimental effects on early postpartum milk production were observed in some studies, particularly in those that restricted energy intake below requirements. Relatively little attention was paid to controlling for adequate protein supply when controlling energy intake.

Cornell research investigated the effects of 3 different dietary energy strategies during the dry period: a bulky, high-fiber controlled energy diet (approximately 100% of energy requirements); a high-energy diet (approximately 150% of energy requirements); and a step-up approach where the controlled energy diet was fed during the first 28 days after dry-off, after which cows were fed an intermediate diet (approximately 125% of energy requirements) for the remainder of the dry period (28 days before expected calving). All diets were fed ad libitum and predicted metabolizable protein (MP) supply was formulated for approximately 1300 g/d. Our observations confirmed that feeding a controlled energy diet prepartum was associated with lower postpartum concentrations of markers of negative energy balance, such as NEFA and BHB, whereas milk production was not different between the groups. In addition, as previously observed by others,<sup>3,10</sup> glucose and insulin concentrations remained higher post-partum in the controlled energy group.<sup>17</sup> This is of great importance for the fresh cow, as glucose is necessary for normal immune cell function and insulin prevents excessive breakdown of adipose and muscle tissue due to its direct inhibitory effects on these processes. Furthermore, high concentrations of BHB and NEFA, as well as lower circulating concentrations of glucose and insulin as observed in cows overfed energy prepartum, have been associated with decreased reproductive success in a number of studies.<sup>3,14,20</sup>

#### *Metabolizable protein (MP) in the dry period*

The drop in dry matter intake around the time of calving as well as the relatively slow increase in intakes in early lactation do not only affect the energy balance of fresh cows, but also their protein balance. Protein and amino acids are instrumental to many physiological functions, particularly cell renewal and immune system function, both of which are particularly important in the transition cow as this is the highest at risk period for infectious diseases like metritis and mastitis. Research shows that protein mobilization starts in the last 2 weeks before calving and carries on until about week 6 post-partum. A majority of mobilized protein is used for milk protein synthesis, with a smaller proportion being used for glucose synthesis. Our current recommendation for an adequate protein supply during the close-up period is 1,200 to 1,400 g/d of predicted MP. Particularly with controlled energy diets, adequate sources of rumen undegradable protein (RUP) should be included in the diets to achieve this goal. No beneficial effects on post-partum performance or health have been observed when higher than recommended amounts of MP were fed. When considering

the cost of protein feed sources and environmental implications of excess nitrogen excretion, feeding protein in great excess of requirements is unwarranted.

#### *Effects of dry period plane of energy on colostrum composition*

Most of the research on the effect of prepartum diet on colostrum composition of cattle stems from research in beef cattle. Few studies have evaluated the effect of feeding dairy cows either a controlled or higher energy diet on colostrum quality and quantity while controlling for adequate protein supply. Our research showed that cows fed a controlled energy diet for the whole duration of the dry period (approximately 57 d) had a greater concentration of IgG in colostrum (96 g/L) than those fed a higher energy diet (72 g/L) during the dry period. At the same time colostrum volume was not significantly different (13 vs 16 lb; 6 vs 7.3 kg).<sup>16</sup> Higher concentrations of IgG in colostrum allow for a higher amount of antibodies to be administered to the calf in 1 feeding, which we consider beneficial for passive transfer of immunity. In our opinion and according to experience shared by others, it is important to allow for an adequate supply of MP prepartum while controlling the diet for energy to prevent a drop in colostrum volume.

#### *Feeding management of dry cow rations*

Even the best formulated rations will not be effective if they are not well-implemented. Bulky rations with the forage base consisting of either straw or mature, low-potassium hay blended with corn silage and a grain mix can be easily sorted by cows if the straw or hay is not chopped, ideally prior to mixing into the TMR. In new research conducted by our group<sup>13</sup> and involving 72 commercial dairy farms in New York and Vermont, only 25% of the prefresh TMR had particle size within recommended ranges (10 to 20% on the top screen; 50 to 60% in the middle; < 40% in the pan) using the Penn State Particle Separator (PSPS). We recommend chopping the straw or hay such that the long particles are no more than 1.5 inches (3.8 cm) (33% on each of the 3 sections of the PSPS). Often, addition of water or another wet ingredient to decrease the ration dry matter into the 46 to 48% range is also required for optimal effectiveness of these rations. Accuracy and consistency in feed delivery and composition are paramount to a successful transition feeding program.

#### *Emerging concepts in feeding the fresh cow*

Ironically, the vast majority of transition cow nutritional management research conducted over the past 20+ years has focused almost exclusively on the dry cow. In most studies focused on transition cow nutrition, dietary treatments were imposed during the prepartum period only and cows were fed a common diet during the post-calving period. Fresh cow rations are common in the dairy industry, although often they are modest variations of the high cow ration, perhaps with slightly higher fiber content and/or the inclusion of modest amounts (1.5 lb (0.7 kg) or less) of straw or hay, lower starch

content, additional rumen undegradable protein, increased amounts of supplemental fat, or targeted inclusion of other nutrients or additives (e.g., rumen-protected choline, additional yeast or yeast culture, additional monensin). Success of these strategies was gauged largely at the farm level, because until recently very few controlled research studies examined these factors in the ration fed during the immediate post-calving period.

#### *Starch and fiber interactions during the pre-calving and post-calving periods*

The research groups at Cornell and the Miner Institute have completed 3 experiments evaluating starch content of the post-partum diet and starch content of the postpartum diet and monensin supplementation throughout the periparturient period.<sup>7,19,24</sup> Dann and Nelson<sup>7</sup> fed 72 multiparous Holstein cows a controlled energy diet during a shortened (40 d) dry period and then 1 of 3 dietary starch regimens during early lactation—a low starch (21.0% starch) diet for the first 91 d postpartum, a medium starch (23.2% starch) diet for the first 21 d post-partum followed by a high starch (25.5% starch) diet through 91 d post-partum, and a high starch diet (25.5% starch) for the first 91 d post-partum. Cows fed the low starch and medium-high starch diets after calving had similar DMI and performance post-calving, whereas cows fed the higher starch diet post-calving had lower DMI and lower milk yield.

McCarthy et al<sup>19</sup> fed primiparous (n = 21) and multiparous (n = 49) Holstein cows diets containing either 26.2% or 21.5% starch from calving through d 21 postpartum; beginning on d 22 postpartum all cows were fed the diet containing 26.2% starch through d 63 postpartum. Cows were also fed either 0 or 400 mg/d of monensin beginning 21 d before expected calving and either 0 or 450 mg/d of monensin beginning at calving and continuing through d 63 post-partum. In contrast to the Miner Institute study, cows fed higher starch diets had faster increases in milk yield and DMI along with lower plasma NEFA and BHBA consistent with better energy status. Cows fed monensin had higher post-partum DMI and milk yield and lower plasma BHBA, regardless of starch level in the post-partum diet.

The Miner and Cornell studies suggest apparently opposite responses to feeding low- and high-starch diets during the fresh period. However, the pre-calving diets were very different between the 2 studies. In the Miner study, cows were fed a typical low starch (13.5% of DM), controlled energy diet for the entire 40-d dry period whereas in the Cornell study, cows were fed a moderate starch close-up diet (17.4% of DM). We speculate that the differences in starch levels between pre-calving and post-calving diets should be no more than 8 to 10 percentage units; cows fed lower starch diets (12 to 14%) immediately before calving should be transitioned onto a fresh diet containing no more than 21 to 22% starch. On the other hand, cows fed higher starch rations before calving (16 to 18% starch) likely can be transitioned onto fresh rations

containing 26 to 27% starch as long as there is sufficient physically effective fiber in the fresh cow diet.

Based upon some case study work as part of a controlled experiment,<sup>19</sup> we also speculate that there are interactions between starch and fiber levels in the post-calving diet. When we had insufficient physically effective fiber in the fresh diet, DMI was higher for cows fed a lower starch diet. However, when straw was increased to levels higher than typical (~ 11% of diet dry matter compared to typical 2 to 4%), DMI was higher for cows fed the higher starch diet. We are currently following up this work with controlled research to further understand the role of fiber in the fresh cow ration.

#### *Additional requirements for metabolizable protein and amino acids in fresh cows?*

In addition to being in negative energy balance, cows also are in negative protein balance during early lactation. This negative protein balance reaches its low point at about 7 days after calving, and cows likely reach positive protein balance by about 21 days after calving.<sup>1</sup> Cows compensate for this negative protein balance in part by mobilizing body protein post-calving, although we understand this process much less than we do the mobilization of body fat during early lactation.

Recently, Larsen et al employed an innovative experimental approach in which they estimated the negative MP balance in cows during the postpartum period and then infused casein into the abomasum in order to eliminate the deficit in MP.<sup>12</sup> Controls received a water infusion, and treatment cows received casein planned to supply 360 g at 1 d postpartum and 720 g at 2 d postpartum, followed by daily reductions of 19.5 g/d ending at 194 g/d at 29 d postpartum. The casein infusion resulted in a high and nearly constant supply of MP from 2 to 29 d postpartum. Although the number of cows in this experiment was very small (n = 4 per treatment), cows infused with casein produced an impressive 7.2 kg/d (~ 16 lb/day) more milk than controls during the experimental period. Further research is needed to evaluate cow responses to supplies of both total MP and individual amino acids during the postpartum period.

### Conclusions

Nutritional management strategies for transition cows have evolved significantly over the past 10 years and continue to evolve. Formulation and implementation of anionic diets pre-calving improves both calcium status and performance (feed intake and milk production) post-calving. Controlled energy diets pre-partum moderate the dynamics of DMI, BCS, and fat mobilization; effective implementation of these diets through excellent feeding management improves metabolic health of fresh cows. Although research focused specifically on nutrition of the fresh cow is limited, new results suggest that there are interactions between starch levels pre-calving and post-calving along with opportunities to combine higher

fermentability with higher effective fiber levels to maintain rumen stability. Furthermore, there appear to be opportunities to focus on metabolizable protein and amino acid nutrition not just during the pre-calving period, but also during the immediate post-calving period.

### References

1. Bell AW, Burhans WS, Overton TR. Protein nutrition in late pregnancy, maternal protein reserves, and lactation performance in dairy cows. *Proc Nutr Soc* 2000; 59:119-126.
2. Block E. Manipulating dietary anions and cations for prepartum dairy cows to reduce incidence of milk fever. *J Dairy Sci* 1984; 67:2919-2948.
3. Cardoso FC, LeBlanc SJ, Murphy MR, Drackley JK. Prepartum nutritional strategy affects reproductive performance in dairy cows. *J Dairy Sci* 2013; 96:5859-5871.
4. Chapinal N, LeBlanc SJ, Carson ME, Leslie KE, Godden S, Capel M, Santos JE, Overton MW, Duffield TW. Herd-level association of serum metabolites in the transition period with disease, milk production, and early lactation reproductive performance. *J Dairy Sci* 2012; 95:5676-5682.
5. Charbonneau E, Pellerin D, Oetzel GR. Impact of lowering dietary cation-anion difference in nonlactating dairy cows: a meta-analysis. *J Dairy Sci* 2006; 89:537-548.
6. Dann HM, Litherland NB, Underwood JP, Bionaz M, D'Angelo A, McFadden JW, Drackley JK. Diets during far-off and close-up dry periods affect periparturient metabolism and lactation in multiparous cows. *J Dairy Sci* 2006; 89:3563-3577.
7. Dann HM, Nelson BH. Early lactation diets for dairy cattle -- focus on starch. In: *Proceedings, Cornell Nutrition Conference for Feed Manufacturers* 2011; 46-56.
8. Drackley JK, Janovick Guretzky NA. Controlled energy diets for dry cows. In: *Proceedings, Western Dairy Management Conference* 2007; 7-16.
9. Gaynor PJ, Mueller FJ, Miller JK, Ramsey N, Goff JP, Horst RL. Parturient hypocalcemia in Jersey cows fed alfalfa haylage-based diets with different cation to anion ratios. *J Dairy Sci* 1989; 72:2525-2531.
10. Janovick NA, Boisclair YR, Drackley JK. Prepartum dietary energy intake affects metabolism and health during the periparturient period in primiparous and multiparous Holstein cows. *J Dairy Sci* 2011; 94:1385-1400.
11. Jardon PW. Using Urine pH to monitor anionic salt programs. *The Compendium for Continuing Education for the Practising Veterinarian* 1995; 17:860-862.
12. Larsen M, Lapierre H, Kristensen NB. Abomasal protein infusion in postpartum transition dairy cows: effect on performance and mammary metabolism. *J Dairy Sci* 2014; 97:5608-5622.
13. Lawton AB, Mann S, Burhans WS, Nydam DV, Rossiter-Burhans CA, Tetreault M, Overton TR. Association of periparturient nutritional strategy with concentration of postpartum  $\beta$ -hydroxybutyrate in dairy cows. *J Dairy Sci* 2015; 98(Suppl. 2):137.
14. Lucy MC. Functional differences in the growth hormone and insulin-like growth factor axis in cattle and pigs: implications for post-partum nutrition and reproduction. *Reproduction in domestic animals = Zuchthygiene* 2008; 43 Suppl 2:31-39.
15. Mann S, Leal Yepes FA, Overton TR, Wakshlag JJ, Lock AL, Ryan CM, Nydam DV. Dry period plane of energy: Effects on feed intake, energy balance, milk production, and composition in transition dairy cows. *J Dairy Sci* 2015; 98:3366-3382.
16. Mann S, Leal Yepes FA, Overton TR, Lock AL, Lamb SV, Wakshlag JJ, Nydam DV. Effect of dry period dietary energy level in dairy cattle on volume, concentration of IgG, insulin and fatty acid composition of colostrum. *J Dairy Sci* 2016; 99:1515-1526.
17. Mann S, Yepes FA, Duplessis M, Wakshlag JJ, Overton TR, Cummings BP, Nydam DV. Dry period plane of energy: Effects on glucose tolerance in transition dairy cows. *J Dairy Sci* 2016; 99:701-717.
18. Martinez N, Risco CA, Lima FS, Bisinotto RS, Greco LF, Ribeiro ES, Maunsell F, Galvao K, Santos JE. Evaluation of periparturient calcium status, energetic profile, and neutrophil function in dairy cows at low or high risk of developing uterine disease. *J Dairy Sci* 2012; 95:7158-7172.

19. McCarthy MM, Dann HM, Overton TR. Feeding the fresh cow. In: *Proceedings*. Cornell Nutrition Conference for Feed Manufacturers 2015.
20. Ospina PA, McArt JA, Overton TR, Stokol T, Nydam DV. Using nonesterified fatty acids and beta-hydroxybutyrate concentrations during the transition period for herd-level monitoring of increased risk of disease and decreased reproductive and milking performance. *Vet Clin North Am Food Anim Pract* 2013; 29:387-412.
21. Reinhardt TA, Lippolis JD, McCluskey BJ, Goff JP, Horst RL. Prevalence of subclinical hypocalcemia in dairy herds. *Vet J* 2011; 188:122-124.
22. Sweeney BM, Ryan CM, Stokol T, Zanzalari K, Kirk D, Overton TR. The effect of decreasing dietary cation-anion difference in the prepartum diet on urine pH and plasma minerals in multiparous Holstein cows. *J Dairy Sci* 2015; 98(Suppl. 2):128.
23. Sweeney BM, Ryan CM, Zanzalari K, Kirk D, Overton TR. The effect of decreasing dietary cation-anion difference in the prepartum diet on dry matter intake, milk production and milk composition in multiparous Holstein cows. *J Dairy Sci* 2015; 98(Suppl. 2):756.
24. Williams SE, Tucker HA, Koba Y, Suzuki R, Dann HM. Effect of dietary starch content on the occurrence of subacute ruminal acidosis (SARA) and inflammation in fresh dairy cows. *J Dairy Sci* 2015; 98(Suppl. 2):741-742.