

Trichomoniasis in Beef Herds

Soren P. Rodning, DVM, MS, DACT
Auburn University, Auburn, AL 36830

Abstract

Reproductive efficiency is one of the most important factors influencing the economic success of a cow-calf operation. Bovine trichomoniasis, also known as trichomonosis, is a venereal disease caused by the protozoan *Tritrichomonas foetus* that negatively impacts a herd's reproductive efficiency, thereby having a substantial economic impact on a cattle operation. This paper outlines the pathogenesis, prevalence, economic impact and diagnosis of trichomoniasis in cattle, as well as reviewing common guidelines for the prevention and control of trichomoniasis.

Résumé

L'efficacité reproductive est l'un des principaux facteurs dont dépend la réussite économique d'un élevage de veaux d'embouche (atelier vache-veau). La trichomonase bovine, aussi appelée trichomonose, est une maladie vénérienne causée par le protozoaire *Tritrichomonas foetus*, qui nuit à la reproduction d'un troupeau et réduit ainsi substantiellement la rentabilité de l'élevage. Cet article décrit la pathogénèse, la prévalence, l'impact économique et le diagnostic de la trichomonase des bovins, tout en rappelant les directives courantes de prévention contre cette maladie.

Introduction

Trichomoniasis is a major cause of fetal wastage that results in substantial economic losses wherever natural breeding conditions exist. Infected bulls are often asymptomatic carriers of *Tritrichomonas foetus* (*T. foetus*); however, they are capable of transmitting the organism to a cow or heifer during coitus. Infections in cows and heifers can result in early embryonic death, abortion, pyometra, fetal maceration, or infertility,^{15,23,40,44} negatively impacting the economic success of a cattle operation. The economic losses associated with trichomoniasis are due to three factors: 1) reduced calf crop due to early embryonic loss or abortion; 2) reduced weaning weight due to delayed conception; and 3) culling and replacement of infected cattle. Since no legal treatment exists in the United States (US), veterinarians and cattle producers must focus on preventive management and other control measures. Understanding the pathogenesis, prevalence, economic impact and

diagnosis of trichomoniasis will assist with implementation of appropriate prevention and control programs.

Pathogenesis and Clinical Signs

In bulls, *T. foetus* localizes in the smegma of the epithelial lining of the penis, prepuce and distal urethra.⁴³ *Tritrichomonas foetus* causes no penile or preputial lesions and does not affect semen quality or libido.^{3,43} An infected bull, therefore, acts only as an asymptomatic carrier, and rarely clears the infection regardless of time. Deep preputial and penile epithelial crypts provide the appropriate microaerophilic environment required for establishment of chronic infections.^{7,11,33,43} A common belief is that *T. foetus* infections in young bulls (less than 3-4 years of age) tend to be transient. However, any bull exposed to *T. foetus* in a natural breeding situation is capable of becoming chronically infected, regardless of age.

Tritrichomonas foetus infection in the cow occurs during coitus with an infected bull. The organism transverse the cervix and colonizes the entire reproductive tract within 1-2 weeks,⁶ and as the organism multiplies in the uterus it can cause death of the embryo or fetus, most commonly between gestational days 15 to 80.⁵¹ A small percentage of cows will not abort until the second or even third trimesters, and an even smaller number of cows (less than 1%) will maintain an infection through a normal gestation and deliver a live calf.⁵⁴ The few cows that maintain a *T. foetus* infection throughout gestation and into the next breeding season are very damaging since they represent a source of reinfection for the herd. Pyometra and abortion are often the first physical signs of trichomoniasis noticed in a herd, but these signs occur in less than 5% of infected animals.³ Infertility due to embryonic death is the most economically damaging symptom and occurs in a larger percentage of infected cows. An affected cow's interestrus interval is usually prolonged because the embryonic loss typically occurs after maternal recognition of pregnancy (days 15-17 of gestation).⁶

Unlike the bull, the cow typically mounts an effective immune response to *T. foetus*,⁵⁶ but the time it takes to clear *T. foetus* from the cow's reproductive tract is quite variable. Primary infections may be cleared from the reproductive tract in as little as 95 days⁴⁴ or as long as 22 months.² Subsequent infections are cleared in about 20 days, indicating an anamnestic response.⁴⁶

Immunity does not persist, however, and the anamnestic response is only significant if reinfection occurs within about 15 months of the primary infection.^{15,43} A cow in a herd with a long breeding season could therefore become pregnant and infected with *T. foetus* early in the breeding season, lose that embryo, be infertile for several months, clear the initial *T. foetus* infection, rebreed, conceive and carry a calf to term as a result of temporary immunity. The result is that more cows will calve later in the calving season than desired, and there is a resultant wide variety in weaning weights rather than just a reduced calving percentage. The later born calves are then marketed at lighter weights, or the cattle producer will incur increased feeding costs to achieve a desired market weight. In either case the cattle producer will sustain substantial economic losses.

Prevalence

Several estimates are available regarding prevalence of trichomoniasis in different regions of North America. In 1964, Johnson reported a 7.5% prevalence in western range bulls.³⁴ More recent studies from Florida,¹ Oklahoma⁶¹ and California⁸ found prevalence rates of 7.3, 7.8 and 4.1%, respectively. The Florida and Oklahoma studies sampled bulls from sale barns or abattoirs, while the California study sampled bulls from randomly selected herds. Rae *et al* conducted an even more recent epidemiological survey of randomly selected natural service beef herds in Florida between 1997 and 1999, and reported a 6% prevalence of *T. foetus*-infected bulls.⁴⁹ Riley *et al* also reported a 6% prevalence in bulls in Saskatchewan, Canada.⁵⁰ In other parts of the world, Erasmus *et al* reported a 7% prevalence in the North Western Cape Province, Western Transvaal and the Orange Free State in South Africa.²⁰

Economic Impact

The economic impact of trichomoniasis is due to: 1) reduced calf crop from early embryonic loss or abortion; 2) reduced weaning weight due to delayed conception; and 3) culling and replacement of infected cattle. Rae developed a computer simulation model to study the impact of trichomoniasis on a cow-calf producer's profitability.⁴⁷ The model estimated a 14 to 50% reduction in annual calf crop if *T. foetus* infections were present in 20 to 40% of the bull population, and the net return per cow exposed to an infected bull decreased by 5 to 35%.⁴⁷ The economic impact of trichomoniasis can be so devastating that several western states in the US consider trichomoniasis a reportable disease and require bull testing prior to sale, prior to transport into the state, or before the use of public land.

Diagnosis of Bovine Trichomoniasis

Diagnosis of *T. foetus* has traditionally relied upon microscopic identification of key morphological characteristics in preputial smegma or cervicovaginal mucus (CVM) incubated in various culture media. Such characteristics include three anterior flagella, one posterior flagellum and an undulating membrane resulting in a jerky movement pattern. However, accurate microscopic identification of *T. foetus* can be complicated by the presence of other trichomonadid protozoa.^{10,16,21,30,58} Contamination of the preputial orifice, prepuce, or penis with fecal material probably explains the presence of these opportunistic trichomonads. Several non-pathogenic protozoa are normal inhabitants of the bovine gastrointestinal tract,^{14,31,39} and therefore proper cleaning of the preputial orifice and proper sampling techniques are critical to avoid fecal contamination of diagnostic samples. None of the contaminating trichomonads, however, results in reproductive pathology in cows or bulls.¹⁷ Therefore, research has recently focused on molecular-based assays to accurately differentiate *T. foetus*^{21,22,32,50} from other trichomonads. Given the lack of legal therapy for bulls infected with *T. foetus*, the only reasonable course of action is to slaughter an infected bull. It is therefore imperative to correctly identify *T. foetus*-infected bulls and not misdiagnose based on the presence of non-pathogenic fecal trichomonads.

At present, molecular-based assays are most commonly used as confirmatory tests for bovine trichomoniasis because of the relatively low cost of *in vitro* cultivation compared to molecular-based assays. However, molecular-based assays are currently very effective in diagnosing human trichomoniasis caused by *Trichomonas vaginalis*, with a sensitivity of 95% and a specificity of 98%.⁴⁵ It is therefore very likely that in the future the preferred diagnostic test for bovine trichomoniasis will be a molecular-based assay, and some researchers have already advocated their use as an independent diagnostic test for bovine trichomoniasis.^{28,42}

Sampling techniques in the male

Several sampling techniques are utilized for obtaining diagnostic specimens in the bull including: 1) a swab technique;⁴¹ 2) a dry pipette technique;^{46,53} 3) a wet pipette technique;³⁶ and 4) the douche technique.³⁶ Fitzgerald *et al* compared the swab and pipette techniques and reported that the number of parasites recovered via the swab technique is only 20% of the number of parasites recovered via pipette scraping.²⁴ The swab technique is therefore rarely used in the US. The dry pipette technique is one of the most common sampling methods in the US, while the douche method is the preferred technique in Europe.⁵³ Schönmann *et al*

reported that the two methods are not statistically different.⁵³

Regardless of technique used, it is generally recommended that bulls be sexually rested 1-2 weeks before testing for *T. foetus*; otherwise, false-negative results are more likely because breeding mechanically removes many of the organisms from a bull's penis and prepuce. Given the sensitivity of *T. foetus* cultures, false-negative results are also possible even if a bull has been sexually rested. Only with three negative tests at weekly intervals (Figure 1) can a veterinarian or producer be 99% sure that a bull is *T. foetus* negative.²⁵

Sampling techniques in the female

Researchers investigating diagnostic sampling methodologies for *T. foetus* have focused primarily on optimizing sample collection and culture from bulls because of their propensity to develop chronic infections. The technique most commonly used to sample female cattle for *T. foetus* is a dry pipette technique.^{7,36,46} An infusion pipette is used to aspirate cervicovaginal mucus (CVM) from the vaginal fornix or near the external cervical os. Alternatively, in the case of a post-coital pyometra, an infusion pipette can also be used to aspirate some of the content of the pyometra. Either sample is then examined directly or placed into appropriate culture medium. Culturing *T. foetus* from cervicovaginal mucus has a reported sensitivity of 58 to 75%.⁵⁵ Samples can also be evaluated with appropriate molecular-based assays.

In vitro culture

Direct microscopic examination of specimens for *T. foetus* is diagnostic, but a far more sensitive method for the detection of *T. foetus* is *in vitro* culture of preputial smegma or CVM in a selective nutrient medium for up to a week.^{35,55,59} *In vitro* culture allows the proliferation of *T. foetus* to more readily detectable levels. All cultures containing organisms resembling *T. foetus* should be confirmed with appropriate molecular-based assays to avoid false-positive results due to fecal trichomonad contamination of culture media.^{10,13,16} Alternatively, samples may be submitted directly for molecular-based evaluation.

	Result	Sensitivity (in series)
First test	Negative	80%
Second test (one week later)	Negative	96%
Third test (one week later)	Negative	99%

Figure 1. Sensitivity (in series) of *T. foetus* cultures.²⁵

In vitro culture media

Various culture and transport media systems have been used including Kupferberg medium and broth, Claussen's medium, Sutherland medium, trypticase-yeast extract-maltose (TYM) medium, Diamond's medium and most recently the InPouch™ TF^a *Tritrichomonas foetus* culture pouch. *In vitro* cultivation using either Diamond's medium or the InPouch™ TF is currently the most common method used to diagnose *T. foetus* in the US. Both culture systems are fairly equal in sensitivity.^{4,7,53,57} However, the InPouch™ TF is somewhat more convenient than Diamond's medium.¹² The InPouch™ TF has a 12-month shelf-life at room temperature, compared to a much shorter refrigerator-life for Diamond's medium. Also, the plastic pouch design of the InPouch™ TF is less likely to break or leak than tubes containing Diamond's medium. Unfortunately, the InPouch™ TF is more expensive than Diamond's medium.

For many years, cultivation of microorganisms with motility and morphology resembling *T. foetus* in either the InPouch™ TF or Diamond's medium was considered to be 100% specific. However, accurate microscopic identification of *T. foetus* has since been shown to be complicated by the presence of other contaminating trichomonadid protozoa.^{10,16,21,30,58} All cultures containing organisms resembling *T. foetus* should therefore be confirmed with appropriate molecular-based assays, or samples should be submitted directly to a laboratory for molecular analysis. Contact the laboratory prior to sample collection to verify the appropriate transport medium.

Treatment

One of the complicating factors associated with bovine trichomoniasis is that there are currently no effective treatments with US Food and Drug Administration approval.⁴⁶ Historically, the most successful treatment for bulls with trichomoniasis involved systemic treatment with nitromidazole derivatives.^{5,19,26,55} However, the use of nitromidazole derivatives is now illegal in food-producing animals in the US because of their mutagenic and carcinogenic properties, and no alternative treatments are available. The lack of effective approved therapies for bovine trichomoniasis emphasizes the need for appropriate preventive and control measures.

Prevention and Control of Bovine Trichomoniasis

Preventing the introduction of *T. foetus* into a cattle herd and controlling trichomoniasis in an infected herd follow many of the same management strategies, and to a large extent focus on herd biosecurity. Ideally, ev-

ery cattle operation should focus on preventing the introduction of *T. foetus*.

Recommended practices to prevent introduction of *T. foetus* into a cattle herd include:

- 1) When possible, avoid grazing cattle on public lands where both bulls and cows have a much greater risk of exposure through coitus with other *T. foetus*-infected animals.²⁷
- 2) Utilize artificial insemination when possible.⁴⁶
- 3) Cull all open cows and heifers.
- 4) Control animal movement into a herd. Maintain good fences to prevent *T. foetus*-infected animals from inadvertently entering a herd, or to prevent uninfected animals from temporarily entering a *T. foetus*-infected herd and then returning with *T. foetus* to their uninfected herd of origin.
- 5) Purchase virgin bulls and heifers as replacements. Buying older bulls and cows as replacements greatly increases the chance of purchasing a *T. foetus*-infected animal. While older bulls are much more likely to become chronically infected with *T. foetus* than cows, a small percentage of cows will also become chronically infected.⁵⁴
- 6) Test bulls for *T. foetus* at least once before introducing them into a new herd.⁴⁶ The test should be performed after two weeks of sexual rest. Ideally, a bull should have three negative cultures at weekly intervals.
- 7) Maintain as young a bull battery as possible. Older bulls are considered more likely to develop chronic *T. foetus* infections.^{43,48} However, any bull exposed to *T. foetus* in a natural breeding situation is capable of becoming chronically infected, regardless of age.
- 8) Breed purchased cows and heifers in a separate herd, and cull all open animals. Ideally, continue to keep the pregnant animals segregated from the rest of the herd through the next breeding season.³⁶
- 9) Consider immunization against *T. foetus* in high-risk herds.

Recommendation for control of trichomoniasis in an infected herd includes:

- 1) Test and cull all infected bulls. Infected bulls should be sold for slaughter only.
- 2) Decrease the number of bulls per breeding unit. Single-sire herds offer the lowest exposure potential. However, single-sire units may not always be practical.
- 3) Reduce the average age of the bull herd. Older bulls are considered more likely to develop chronic *T. foetus* infections.^{43,48} However, any bull

exposed to *T. foetus* in a natural breeding situation is capable of becoming chronically infected, regardless of age.

- 4) Test bulls for *T. foetus* at least once before introducing them into a new herd.⁴⁶ The test should be performed after two weeks of sexual rest. Ideally, a bull should have three negative cultures at weekly intervals.
- 5) Utilize artificial insemination when possible.⁴⁶
- 6) Reduce the breeding season to 60-90 days and cull all open cows and heifers. If there are too many open cows for culling to be economically feasible, then at least these animals should be separated into a high-risk herd. A long breeding season not only allows propagation of *T. foetus*, but it may also hide production losses due to reduced weaning weights because of delayed conception.³⁷
- 7) Culture all pyometras diagnosed in cows or heifers during pregnancy examinations.
- 8) Submit all aborted fetuses and placental tissue to a diagnostic laboratory.
- 9) Immunization against *T. foetus* is an extremely important management tool for herds infected with *T. foetus*. Research trials clearly demonstrate the benefit of *T. foetus* vaccination.^{9,18,29,38,52,60} TrichGuard^{®b} and TrichGuard[®] V5L^c are currently the only *T. foetus* vaccines available in the United States. The vaccines require an initial subcutaneous dose followed by a booster dose two to four weeks later. The second injection should precede the breeding season by four weeks. Annual revaccination four weeks prior to the breeding season is recommended.

Endnotes

^a InPouch™ TF *Trichomonas foetus* culture pouch – BioMed Diagnostics, White City, OR

^b TrichGuard[®] – Fort Dodge Animal Health, Fort Dodge, IA

^c TrichGuard[®] V5L – Fort Dodge Animal Health, Fort Dodge, IA

References

1. Abbitt B, Meyerholz GW: *Trichomonas foetus* infection of range bulls in South Florida. *Vet Med Sm Anim Clin* 74:1339-1342, 1979.
2. Alexander GI: An outbreak of bovine trichomoniasis in Queensland and its control. *Aust Vet J* 29:61-69, 1953.
3. Anderson ML, Barr BC, Conrad PA: Protozoal causes of reproductive failure in domestic ruminants. *Vet Clin North Am Food Anim Pract* 10:439-461, 1994.
4. Appell LH, Mickelsen WD, Thomas MW, Harmon WM: A comparison of techniques used for the diagnosis of *Trichomonas foetus* infections in beef bulls. *Agri Pract* 14:30-34, 1993.

5. Ball L, Dargatz DA, Cheney JM, Mortimer RG: Control of venereal disease in infected herds. *Vet Clin North Am Food Anim Pract* 3:561-574, 1987.
6. BonDurant RH: Diagnosis, treatment, and control of bovine trichomoniasis. *Compend Contin Educ Pract Vet* 7:S179-188, 1985.
7. BonDurant RH: Pathogenesis, diagnosis and management of trichomoniasis in cattle. *Vet Clin North Am Food Anim Pract* 13:345-361, 1997.
8. BonDurant RH, Anderson ML, Blanchard P, et al: Prevalence of trichomoniasis among California beef herds. *J Am Vet Med Assoc* 196:1590-1593, 1990.
9. BonDurant RH, Corbeil RR, Corbeil LB: Immunization of virgin cows with surface antigen TF1.17 of *Tritrichomonas foetus*. *Infect Immun* 61:1385-1394, 1993.
10. BonDurant RH, Gajadhar A, Campero CM, et al: Preliminary characterization of a *Tritrichomonas foetus*-like protozoan isolated from preputial smegma of virgin bulls. *Bov Pract* 33:124-127, 1999.
11. BonDurant RH, Honigberg BN: Trichomonads of veterinary importance, in Kreier JP (ed): *Parasitic Protozoa*, ed 2. San Diego, Academic Press, 1994, pp 111-188.
12. Borchardt KA, Norman BB, Thomas MW, Harmon WM: Evaluation of a new culture method for diagnosing *Tritrichomonas foetus* infection. *Vet Med* 87:104-112, 1992.
13. Campero CM, Rodriguez Dubra C, Bolondi A, et al: Two-step (culture and PCR) diagnostic approach for differentiation of non-*T. foetus* trichomonads from genitalia of virgin beef bulls in Argentina. *Vet Parasitol* 112:167-175, 2003.
14. Castella J, Munoz E, Ferrer D, Gutierrez JF: Isolation of the trichomonad *Tetratrichomonas buttrei* (Hibler et al, 1960) Honigberg, 1963 in bovine diarrhoeic faeces. *Vet Parasitol* 70:41-45, 1997.
15. Clark BL, Dufty JH, Parsonson IM: The effect of *Tritrichomonas foetus* on calving rates in beef cattle. *Aust Vet J* 60:71-74, 1983.
16. Cobo ER, Campero CM, Mariante RM, Benchimol M: Ultrastructural study of a tetratrichomonad species isolated from preputial smegma of virgin bulls. *Vet Parasitol* 117:195-211, 2003.
17. Cobo ER, Canton G, Morrell E, Cano D, Campero CM: Failure to establish infection with *Tetratrichomonas* sp in the reproductive tracts of heifers and bulls. *Vet Parasitol* 120:145-150, 2004.
18. Cobo ER, Morsella C, Cano D, Cipolla A, Campero CM: Immunization in heifers with dual vaccines containing *Tritrichomonas foetus* and *Campylobacter fetus* antigens using systemic and mucosal routes. *Therio* 62:1367-1382, 2004.
19. Dawson LJ: Diagnosis, prevention, and control of campylobacteriosis and trichomoniasis. *Bov Pract* 21:180, 1986.
20. Erasmus JA, De Wet JAL, Van der Merwe HE, Pienaar GCJ: Bovine trichomoniasis in the North Western Cape Province, Western Transvaal, and the Orange Free State. *J S Afr Vet Assoc* 60:51-52, 1989.
21. Felleisen RSJ, Lambelet N, Bachmann P, et al: Detection of *Tritrichomonas foetus* by PCR and DNA enzyme immunoassay based on rRNA gene unit sequences. *J Clin Microbiol* 36:513-519, 1998.
22. Felleisen RSJ, Schmid-Lambelet LN, Walubengo J: Comparative evaluation of methods for the diagnosis of bovine *Tritrichomonas foetus* infection. *J Protozool Res* 7:90-101, 1997.
23. Fitzgerald PR: Bovine trichomoniasis. *Vet Clin North Am Food Anim Pract* 2:277-282, 1986.
24. Fitzgerald PR, Hammond DM, Miner MC, Binns W: Relative efficacy of various methods of obtaining preputial samples for diagnosis of trichomoniasis in bulls. *Am J Vet Res* 13:452-457, 1952.
25. Food Animal Reproduction and Herd Health Services, University of California at Davis; Bovine Trichomonosis: Essential Facts and Testing. www.vetmed.ucdavis.edu/vetext/INF-BE_Bov-trichomonosis.html, accessed May 2007.
26. Gasparini G, Vaghi M, Tardani A: Treatment of bovine trichomoniasis with metronidazole (8823 R.P.). *Vet Rec* 75:940, 1963.
27. Gay JM, Ebel ED, Kearley WP: Commingled grazing as a risk factor for trichomonosis in beef herds. *J Am Vet Med Assoc* 209:643-646, 1996.
28. Grahn RA, BonDurant RH, Van Hoosear KA, Walker RL, Lyons LA: An improved molecular assay for *Tritrichomonas foetus*. *Vet Parasitol* 127:33-41, 2005.
29. Hall MR, Kvasnicka WG, Hanks D, Chavez L, Sandblom D: Improved control of trichomoniasis with *Trichomonas foetus* vaccine. *Agri Pract* 14:29-34, 1993.
30. Hayes DC, Anderson RR, Walker RL: Identification of trichomonadid protozoa from the bovine preputial cavity by polymerase chain reaction and restriction fragment length polymorphism typing. *J Vet Diagn Invest* 15:390-394, 2003.
31. Hibler CP, Hammond DM, Caskey FH, Johnson AE, Fitzgerald PR: The morphology and incidence of the trichomonads of swine, *Tritrichomonas suis* (Gruby & Delafond), *Tritrichomonas rotunda* n.sp. and *Trichomonas buttrei* n.sp. *J Protozool* 7:159-171, 1960.
32. Ho MSY, Conrad PA, Conrad PJ, et al: Detection of bovine trichomoniasis with a specific DNA probe and PCR amplification system. *J Clin Microbiol* 32:98-104, 1994.
33. Honigberg BN: Trichomonads of veterinary importance: *Tritrichomonas foetus*, in Kreier JP (ed): *Parasitic Protozoa*, ed 1. New York, Academic Press, 1978, pp 207-273.
34. Johnson AE: Incidence and diagnosis of trichomoniasis in western beef bulls. *J Am Vet Med Assoc* 145:1007-1010, 1964.
35. Johnson AE: The diagnosis of trichomoniasis in the bull. *Proc of the US Livestock Sanitary Assoc* 69:183-189, 1965.
36. Kimsey PB: Bovine trichomoniasis, in Morrow DA (ed): *Current Therapy in Therio*, ed 2. Philadelphia, WB Saunders Company, 1986, pp 275-279.
37. Kvasnicka WG, Hall MR, Hanks DR: Bovine Trichomoniasis, in Howard JL, Smith RA, (ed): *Current Veterinary Therapy 4: Food Animal Practice*, Philadelphia, WB Saunders Co., 1999, pp 420-425.
38. Kvasnicka WG, Hanks D, Huang J, et al: Clinical evaluation of the efficacy of inoculating cattle with a vaccine containing *Tritrichomonas foetus*. *Am J Vet Res* 53:2023-2027, 1992.
39. Levine NP: *Veterinary Protozoology*. Ames, Iowa State University Press, 1985.
40. Mickelsen WD, Paisley LG, Anderson PB: Survey of the prevalence and types of infertility in beef cows and heifers. *J Am Vet Med Assoc* 189:51-54, 1986.
41. Morgan BB: *Bovine Trichomoniasis*. Minneapolis, Burgess Publishing Company, 1946.
42. Nickel DD, Olson ME, Schultz GA: An improved polymerase chain reaction assay for the detection of *Tritrichomonas foetus* in cattle. *Can Vet J* 43:213-216, 2002.
43. Parsonson IM, Clark BL, Dufty JH: The pathogenesis of *Tritrichomonas foetus* infection in the bull. *Aust Vet J* 50:421-423, 1974.
44. Parsonson IM, Clark BL, Dufty JH: Early pathogenesis and pathology of *Tritrichomonas foetus* infection in virgin heifers. *J Comp Pathol* 86:59-66, 1976.
45. Patel SR, Wiese W, Patel SC, et al: Systematic review of diagnostic tests for vaginal Trichomoniasis. *Infect Dis Obstet Gynecol* 8:248-257, 2000.
46. Peter D: Bovine venereal diseases, in Youngquist RS (ed): *Current Therapy in Large Animal Therio*, ed 1. Philadelphia, WB Saunders Company, 1997, pp 355-363.
47. Rae DO: Impact of trichomoniasis on the cow-calf producer's profitability. *J Am Vet Med Assoc* 194:771-775, 1989.
48. Rae DO, Chenoweth PJ, Genho PC, et al: Prevalence of *Tritrichomonas fetus* in a bull population and effect on production in a large cow-calf enterprise. *J Am Vet Med Assoc* 214:1051-1055, 1999.
49. Rae DO, Crews JE, Greiner EC, Donovan GA: Epidemiology of *Tritrichomonas foetus* in beef bull populations in Florida. *Therio* 61:605-618, 2004.
50. Riley DE, Wagner B, Polley L, Krieger JN: PCR-based study of conserved and variable DNA sequences of *Tritrichomonas foetus* isolates from Saskatchewan, Canada. *J Clin Microbiol* 33:1308-1313, 1995.
51. Roberts SJ: Infectious diseases causing infertility in cows, in *Veterinary Obstetrics and Genital Disease*, ed 3. Ann Arbor, MI, Edward Bros, Inc, 1986, p 447.

52. Schnackel JA, Wallace BL, Kvasnicka WG, Hanks DR, Hall MR: *Trichomonas foetus* vaccine immunogenicity trial. *Agri Pract* 10:11-14, 1989.
53. Schönmann MJ, BonDurant RH, Gardner IA, *et al*: Comparison of sampling and culture methods for the diagnosis of *Tritrichomonas foetus* infection in bulls. *Vet Rec* 134:620-622, 1994.
54. Skirrow S: Identification of trichomonad-carrier cows. *J Am Vet Med Assoc* 191:553-554, 1987.
55. Skirrow S, BonDurant RH: Bovine trichomoniasis. *Vet Bull* 58:592-603, 1988.
56. Skirrow SZ, BonDurant RH: Immunoglobulin isotype of specific antibodies in reproductive tract secretions and sera in *Tritrichomonas foetus*-infected heifers. *Am J Vet Res* 51:645-653, 1990.
57. Skirrow SZ, BonDurant RH, Farley J, Correa J: Efficacy of ipronidazole against trichomoniasis in beef bulls. *J Am Vet Med Assoc* 187:405-407, 1985.
58. Taylor MA, Marshall RN, Stack M: Morphological differentiation of *Tritrichomonas foetus* from other protozoa of the bovine reproductive tract. *Brit Vet J* 150:73-80, 1994.
59. Tedesco LF, Errico F, Baglivi LP: Diagnosis of *Tritrichomonas foetus* infection in bulls using two sampling methods and a transport medium. *Aust Vet J* 55:322-324, 1979.
60. Villaruel A, Carpenter TE, BonDurant RH: Development of a simulation model to evaluate the effect of vaccination against *Tritrichomonas foetus* on reproductive efficiency in beef herds. *Am J Vet Res* 65:770-775, 2004.
61. Wilson SK, Kocan AA, Gaudy ET, Goodwin D: The prevalence of trichomoniasis in Oklahoma beef bulls. *Bov Pract* 14:109-110, 1979.