

culosis, Medline, reference lists and hand searching proceedings. A total of 66 research trials were identified, and 20 met the criteria for inclusion in the systematic review.

Results

The JD vaccine reduced the occurrence of clinical disease by 92% (95% CI 90-96%), with a heterogeneity chi square of $p=.96$. The vaccine reduced fecal shedding by 57% (95% CI 41-69%), there was heterogeneity (p

value $=.001$), but all trials were able to reduce fecal shedding. The JD vaccine reduced the numbers of animals with positive tissues by 56% (95% CI 49- 62%). Again, there was heterogeneity (p value $=.000$), but all trials had a reduction in positive tissues.

Significance

These data would suggest that the Johne's disease vaccine is effective at reducing clinical disease, fecal shedding and tissue infection.

Effect of Sub-Clinical Manifestations of Bovine Paratuberculosis in Beef Cattle

Mario A. Villarino, DVM, PhD¹, S.E. Wikse, DVM, ACVP² and E.R. Jordan, PhD¹

¹Texas Cooperative Extension, The Texas A&M University System, Dallas, TX 75252

²College of Veterinary Medicine, Texas A&M University, College Station, TX 77843

Introduction

Bovine paratuberculosis (Johne's disease) is a chronic, debilitating disease of cattle. Although widely known in the dairy industry, the effect of the disease in extensive ranch operations is unknown. Currently, testing techniques for bovine paratuberculosis in beef cattle are few, costly and somewhat inaccurate. We conducted the study of a beef herd located in Texas with the objective of evaluating the effect of subclinical manifestations of bovine paratuberculosis in beef cattle.

Materials and Methods

We conducted serial testing using ELISA for serum samples collected during three consecutive years (2003-2004-2005) of 695 *Bos taurus* x *Bos indicus* cows from a ranch known to have clinical cases of the disease with fecal culture positive samples. Animals were considered positive when an ELISA S/P value of 0.25 or more was detected in at least one of the tested years. Weaning weight and age of wean of the offspring (females, males and steers) of seropositive and seronegative dams were compared using One-Way ANOVA, all pairwise multiple comparison (Tukey test), Dunns'

method of multiple comparison and Kruskal-Wallis one-way analysis of variance in non-parametric comparisons.

Results

The most important finding was that the weaning weights of female offspring of seropositive dams were significantly different than the weaning weights of female offspring of seronegative dams ($P=0.035$). We also found statistically significant differences in weaning weights between females, males and steers in each seropositive and seronegative group (the male offspring of seronegative dam were considerably heavier). No statistically significant differences were found in age at weaning.

Significance

Our results indicate that subclinical cases of bovine paratuberculosis can cause significant differences in herd performance (in our study, weaning weights) on beef herds. Due to the chronicity of the disease and limited performance of the currently available tests, serial testing is required to provide an estimate of disease prevalence.