

Neosporosis in Cattle

Mark Anderson

California Veterinary Diagnostic Laboratory, School of Veterinary Medicine, UC Davis, Davis California

Introduction

Neosporosis is a newly recognized protozoal infection that has a worldwide distribution and is reported to be a major cause of abortion in North America, Europe and New Zealand. An important feature of the disease is that the parasite can be maintained in cattle as a chronic, apparently lifelong, infection which can be passed on to the fetus during pregnancy. In some cows this fetal infection may result in an abortion. However, most infected cows give birth to a healthy, but congenitally infected calf. A heifer calf that is born congenitally infected is capable of transmitting the infection on to the next generation when she becomes pregnant, thus maintaining the infection in the herd. Diagnosis of the infection is assisted through the examination of aborted fetuses and serologic testing of cattle.

This paper is intended to provide the bovine practitioner with current information about neosporosis in cattle. Several review papers and book chapters are available for additional information.^{1,3,23,30,55}

The protozoan parasite *Neospora caninum* was first identified in dogs with encephalomyelitis and myositis.^{13,14,21,22} It has been associated with disease in various species of livestock including cattle, sheep, goats and horses. In cattle, the parasite has been isolated from aborted fetuses and congenitally infected calves.^{10,23} The pathogenic potential of these isolates has been confirmed by experimental infection of pregnant cattle resulting in fetal death and/or congenitally infected calves.^{19,23} These isolates have been used in the development of a variety of serodiagnostic tests.^{12,16,20,23,24,29,37,39,41,45,58,61}

Bovine neosporosis has emerged as an important reproductive disease since its first association with an abortion storm in 1987 on a dairy in New Mexico.⁴⁶ Numerous reports of *Neospora* abortion have confirmed this infection as a significant cause of abortion, particularly among dairy cattle.^{2,7,18,23,25,26,36,45,46,47,48,55} It has a worldwide distribution, having been diagnosed in many countries from six continents, and is prevalent throughout North America. Bovine neosporosis is probably not a new disease, but rather a newly recognized one. Retrospective studies in California have confirmed that the parasite has been endemic since 1984, and a decade earlier the infection was identified retrospectively in a stillborn

calf in Australia.²³ Surveys in California, the Netherlands and New Zealand indicate that approximately 20% of all aborted bovine fetuses submitted to the diagnostic laboratory are diagnosed with this infection. In abortion submissions from dairy herds with a history of *Neospora* abortion, the proportion of *Neospora* infection in aborted fetuses is as much as 44%.⁴ Since fetal infection can result in either fetal death or the birth of a live congenitally infected calf, there are questions about what factors determine the clinical outcome of fetal infection and what are the appropriate methods to diagnose *Neospora* abortion. This will be discussed.

Clinical Presentation

There are no clinical signs in cows that abort due to *Neospora* infection. The aborted fetuses are usually autolyzed with no gross lesions and placentas are not retained. Abortions have been diagnosed in both heifers and cows from three months' gestation to term, but the majority of *Neospora* abortions occur in the second trimester of pregnancy. This pattern of mid-gestation abortion is distinctive from other diagnosed causes of infectious abortion in dairy cattle which tend to occur later in gestation. Whether *Neospora* infection can cause reproductive problems in the first trimester of gestation is unknown. Fetal mummification has been associated with *Neospora* outbreaks. *Neospora* abortions occur throughout the year.

Neospora infections associated with abortion and congenital infections have been reported in both dairy and beef cattle. There are more reports attributing significant numbers of abortions in dairy cattle, particularly those in dry-lot dairies. The reasons for this are not certain. It is likely that the mid-gestation fetuses typical of *Neospora* abortion would more easily be found and submitted for diagnosis in the dry-lot dairy environment. Alternatively, it may be that the environment of the dry-lot dairy is more conducive to the spread and transmission of this disease. The serologic evidence of *Neospora caninum* infection in dairy herds in the U.S. ranges from 5% to 98%. Overall, the seroprevalence in the U.S. is estimated to be approximately 45% in dairy cattle and about 5% in beef cattle, based on submissions to the California Veterinary Diagnostic Laboratory

(Hietala, personal communication 1999). The estimated seroprevalence in dairy cattle in England and Wales is 6% (Trees, Vet. Rec. letter, 1998) and a similar level of seroprevalence was identified in New Zealand cattle (Reichel, N. Z. Vet. J, letter, 1998). A survey of Canada beef cattle found a seroprevalence of 30%.⁵⁷

Cattle with serologic evidence of infection have an increased risk of abortion.^{35,43,52,57,60} In a California study, seropositive congenitally infected cows had a 7.4 increased risk of abortion in their first pregnancy. The risk in the second pregnancy was considerably lower, though this may have been influenced to some degree by selective culling of aborting cows from the first pregnancy.⁵² In the Netherlands, a 3-fold increase in abortion risk was associated with seropositive cows and compared to seronegative herd mates.⁶⁰ In addition to abortion and congenital infection, *Neospora* infection may cause reduced milk production and shortened production life based on a study of seropositive cows which produced less milk and were culled earlier than seronegative herd mates.^{50,51} Seropositive beef cattle have an increased risk of abortion and stillbirth, in addition to increased risk of culling for any reason and culling for poor reproductive performance.⁵⁷

Two patterns, endemic abortion and epidemic abortion, have been described in association with neosporosis in herds of cattle.⁵⁵ In the endemic pattern of abortion, the herd experiences an elevated abortion rate of greater than 5% per year which persists for years. In investigations of two California dairies with endemic *Neospora* abortions, the annual abortion rate attributable to neosporosis in these herds was estimated to be 10.6% and 17.3%.⁵³ The epidemic pattern of abortion is less common and is characterized by abortions in a high proportion of pregnant cattle over a relatively brief period of time. In some instances, over 30% of pregnant cattle have aborted due to neosporosis within several months.⁴⁸ An apparent mixture of these patterns may be observed in some herds that have experienced a prolonged history of sporadic cases of *Neospora* abortion and occasional outbreaks of abortions attributable to *Neospora*.

In most instances, cows that abort a *Neospora*-infected fetus will have either additional abortions,^{4,28} or infected fetuses in subsequent pregnancies.^{9,23} The clinical outcome of these subsequent pregnancies is variable but a seropositive cow that aborts has an estimated 2-fold to 5.7-fold greater risk of abortion in subsequent pregnancies.^{35,52}

An uncommon manifestation of fetal *Neospora* infection is the birth of a clinically affected full-term calf which exhibits variable CNS signs manifested as limb dysfunctions, ranging from mild proprioceptive defects to complete paralysis. Microscopically there is a multifocal protozoal encephalomyelitis which may be particularly localized in the spinal cord gray matter.^{9,23}

The majority of calves that acquire a *Neospora* infection during gestation are born clinically normal. These calves will have a high precolostral antibody titer to *Neospora caninum* which is useful in detecting *in utero* infection. A high percentage, 80% to over 90%, of calves born to seropositive cows are congenitally infected based on serology.^{9,42,45,55} As will be discussed in the section on transmission, these clinically normal, congenitally infected calves are important in maintaining the infection in the herd.

Diagnosis

The confirmation of *Neospora* infection as the cause of abortion will require the assistance of a veterinary diagnostic laboratory. The preferred samples in cases of abortion include 1 or more aborted fetuses submitted with placenta and serum from the dam. The aborted fetus is usually autolyzed with serosanguinous fluid accumulation in body cavities. Rarely there are subtle gross lesions, consisting of pale white foci in the skeletal muscles or the heart. Histologic lesions consist of widespread nonsuppurative infiltrates. The most diagnostically significant lesions are found in the brain and consist of scattered foci of nonsuppurative cellular infiltrates with occasional foci of necrosis. Protozoa are not usually seen on routinely stained slides. Other histologic lesions that are consistently found include nonsuppurative epicarditis and/or myocarditis, focal nonsuppurative myositis and nonsuppurative portal hepatitis, frequently with focal hepatic necrosis and focal nonsuppurative interstitial pneumonia.^{3,7}

Immunohistochemistry using antibodies raised against *Neospora caninum* antigens is an effective method employed by many diagnostic laboratories to identify the tachyzoite and tissue cyst stages of the parasite in fetal tissues. *Neospora* immunohistochemistry is most successful on sections of fetal brain, although the parasites also are frequently present in the lung, kidney and skeletal muscle. Immunohistochemistry has been successfully employed to diagnose *Neospora* infections in mummified fetuses, although the autolytic state of these fetuses diminishes the diagnostic accuracy.

The use of pathology and immunohistochemistry on aborted fetuses to establish a diagnosis of *N. caninum* as the cause of an abortion has been questioned since the fetus can be infected and not abort due to the infection.⁵⁴ The answers to what determines whether a fetus will live or die and how to accurately diagnose the cause of abortion requires an understanding of the parasite and fetal ontogeny. *Neospora caninum* is a relatively weak pathogen that is adapted to cattle and utilizes vertical transmission: in the birth of a live infected calf, rather than abortion, helps maintain the infection in the herd.

Immune-competent animals or fetuses that become infected are likely to develop subclinical infection or overcome the infection. In cattle, fetal immune competence is first detectable at about 120 to 150 days gestation. Fetuses that acquire the infection prior to sufficient development of an immune response become overwhelmed with disseminated infection, resulting in death. These deaths occur in fetuses primarily through the sixth month of gestation, after which the number of *Neospora caninum*-induced fetal deaths is greatly reduced. These disseminated inflammatory lesions occur in the brain, lungs, heart, liver, kidney, muscles, placenta and other organs. Fetal death is presumed to be primarily related to the placental and heart lesions. A lack of fetal resistance to disseminated infection is responsible for the predominance of abortions and fetal mummification that occurs in mid-gestation.

To increase the likelihood of an accurate diagnosis of *Neospora* abortion, the diagnostician must take into consideration the following factors prior to establishing the infection as the cause: gestational age and postmortem condition (autolyzed); presence of compatible disseminated inflammatory lesions; presence of detectable parasites with immunohistochemistry; and lack of other abortifacients. Conversely, a *Neospora caninum*-infected aborted fetus with mild focal lesions (usually consisting of focal encephalitis in late-term fetuses) may have an incidental *Neospora* infection, and other causes for the abortion should be investigated.

A variety of serologic tests are available to assist in the diagnosis of neosporosis. These include the indirect fluorescent antibody test (IFA), the modified agglutination test, and a number of enzyme-linked immunosorbent assays (ELISA).^{12,16,20,24,26,27,28,29,37,39,41,58,61} The assays utilize *Neospora caninum* tachyzoites or specific derived antigens. The specificity and sensitivity of the various serologic tests are comparable, depending on the minimum antibody titer that has been established as the cut-off for a positive result. Laboratories utilizing any of the serologic tests for *Neospora* should establish appropriate cut-off titers using sera from known infected and noninfected cattle. In some tests, the positive cut-off titer has been selected based on the antibody titer in a cow that has aborted an infected fetus. Thus, this cut-off may not be the most appropriate for the serologic diagnosis of a chronic infection in cattle which vary in age and pregnancy status.⁴⁵

A single serum sample from an individual cow may not accurately reflect her infection status, since titers in known positive cattle fluctuate and may fall below the cut-off value for some period of time. In rare instances, cows that abort a *Neospora*-infected fetus may not have a significantly elevated titer. Also, elevated titers at the time of abortion may decline over several months following abortion.²⁰ A portion of congenitally

infected heifers with a history of positive titers, which had infected their fetuses, had titers below the cut-off at the time of parturition.⁵ There is no conclusive evidence to demonstrate that a seropositive cow can revert to a consistently seronegative status.

Neospora serology is effective in detecting elevated *Neospora* antibodies in the serum of congenitally infected or *in utero*-exposed calves.^{9,23} In addition, serology may be useful in establishing the diagnosis in aborted fetuses, since infected fetuses 6 months or more in gestation may have elevated *Neospora* antibody titers. Recent work suggests that the *Neospora caninum* IFA test is the most accurate method to detect fetal antibodies.^{11,59} However, a negative fetal *Neospora* IFA titer does not rule out the possibility of infection, and a positive titer does not prove that this infection was the cause of the abortion.

An ELISA for detection of *Neospora* antibodies has been developed and is used for routine diagnostic testing at the California Veterinary Diagnostic Laboratory as part of the bovine abortion serologic panel.^{41,55} This procedure is rapid, inexpensive and has excellent sensitivity (88.6%) and specificity (96.5%). Cutoff values have been established for the ELISA by which the probability of infection can be estimated in cattle. In the individual aborting cow, a positive serology result does not prove that the abortion was due to neosporosis, but it can assist the diagnosis. In addition to its use in the routine abortion screen in individual abortion cases, the ELISA test is used on a herd basis. The test can be used to estimate herd seroprevalence of *Neospora* infection, investigate the association between seropositivity and abortion, and estimate the extent of herd infection attributable to congenital infection. It isn't clear whether the titer in the individual cow can be used to predict the outcome of an ongoing pregnancy, but cows with high antibody levels at 180 and 210 days' gestation were less likely to abort than cows with low antibody titers at these times.⁴³

Transmission

The ways cattle acquire *Neospora* infection are under investigation. Forms of the parasite that have been identified in infected fetuses and calves are the tachyzoite and tissue cyst stages. Tachyzoites can spread through the body and invade the cells of a variety of organs. The tachyzoite stage is associated with inflammation and necrosis at the site of invasion. The tissue cyst stage, containing multiple bradyzoites surrounded by a thick cyst wall, is found in neural tissues. The tissue cyst elicits minimal inflammatory reaction and can persist for long periods of time.

There are several ways cattle may acquire *Neospora* infection, either by horizontal (postnatal) infection or by vertical transmission of the infection transplacentally during pregnancy. The horizontal

transmission model is based on the fact that *Neospora* is similar and related to other apicomplexan coccidia, such as *Toxoplasma gondii* and *Sarcocystis* spp. The life cycle of this family of protozoa parasites requires two hosts: a carnivorous predator is the definitive host and a prey species is the intermediate host. For example, in toxoplasmosis, the cat is the definitive host and can acquire the infection through ingestion of parasites contained in the prey species. *Toxoplasma gondii* parasites undergo sexual replication in the intestine of the cat and oocysts are shed in the feces. These *Toxoplasma gondii* oocysts are capable of infecting a wide variety of animal species (intermediate hosts). Among livestock species, toxoplasmosis is a significant cause of abortion in sheep and goats, though not in cattle.

The taxonomic and morphologic similarities between *Neospora* and *Toxoplasma* support the hypothesis that the cow may be infected through oral ingestion of coccidial oocysts shed from a carnivorous definitive host. The identification of this proposed definitive host has been difficult. Previous studies on dogs, cats, rats, mice, raccoons and various bird species had confirmed no role for these species in transmission of *Neospora* to cattle.^{6,23} However, recently McAllister, et al were successful in transmitting *Neospora* infection from experimentally infected mice to puppies, which shed oocysts in their feces.³⁴ Experimental transmission of the infection from the dog to cattle has not yet been reported, but the results of this study strongly suggest that the dog is a definitive host. There is also epidemiological evidence which suggests that dogs may play a role in the transmission of neosporosis. Paré, et al found an association between the herd seroprevalence of *Neospora* infection and the presence and number of dogs at the herd.⁴⁴

Although there are close similarities between *Neospora* and *Toxoplasma*, there are differences between neosporosis in cattle and toxoplasmosis in sheep. In sheep toxoplasmosis, when the infection is acquired during pregnancy, the ewe seroconverts to the parasite and fetal infection and/or abortion may occur. In subsequent pregnancies the ewe is resistant to infection. However, neosporosis in pregnant cattle differs because the cow does not need to acquire *Neospora* infection or seroconvert during pregnancy for her fetus to become infected. Unlike toxoplasmosis, cows that abort in one pregnancy are susceptible to repeat infection and are at an increased risk of repeat abortion.

Vertical transmission of *Neospora* through generations of cattle appears to be the major method by which *Neospora* infection is maintained in herds.^{5,15,40,42,43,45,53,55,60} In a Swedish study, the role of congenital transmission of neosporosis was supported by evidence of the familial distribution of seropositive cattle through successive generations. In a German study, 93% of the descendants of seropositive cows were also seropositive, indicating that

vertical transmission was the major method of transmission of infection in the herds examined.⁴⁵ In California dairies, several serologic studies also offer evidence of vertical transmission. In endemic herds, the majority of calves born to seropositive cows have serologic evidence of congenital infection. In addition, the rate of seropositivity in the herd is not associated with the age of the cow, suggestive that the rate of acquired infection after birth is low.

There is additional pathologic and serologic evidence which indicates that these congenitally infected calves have a chronic persistent infection which can be passed on transplacentally to their offspring. In a survey of heifer calves in a known *Neospora* dairy herd, heifer calves with serologic evidence of congenital exposure were compared with serologically negative cohorts. The two groups were similar until calving, at which time all the offspring of the seropositive heifers had elevated *Neospora* titers, all seronegative heifers had serologically negative calves, and there was no evidence of seroconversion to *Neospora* among the negative heifers.⁵ A portion of the calves were necropsied, and the calves from seropositive heifers had histologic lesions in the brain and spinal cord consistent with congenital *Neospora* infection. Protozoa were identified by immunohistochemistry. The calves from seronegative heifers had no lesions or other findings suggestive of congenital *Neospora* infection.

While vertical transmission appears to be the major way that cattle become infected with *Neospora* in endemic herds, there is serologic evidence that cows that have aborted during an epidemic probably acquired the infection after birth based on analysis of the seropositivity of dams and daughters.⁵³ In addition, the pattern of abortion outbreaks in epidemic neosporosis is suggestive of a point-source exposure with acquired infection.^{23,32,36,48} However, there has been no confirmation that the cattle aborting in an epidemic acquired the infection during their pregnancy. In endemically infected herds which have been sampled more extensively, there is serologic evidence that a low level of postnatal infection from unknown sources apparently does occur.^{45,55}

Control and Prevention

A major method of *Neospora* transmission in herds is through the infection of fetuses in cattle that are chronically infected. These infected cows can be identified based on their serologic titers or from a history of *Neospora* abortion or congenital infection. With this knowledge, control of the infection could be focused on reducing the number of infected cows in the herd and limiting the introduction of infected replacement cattle into the herd. Culling decisions concerning cows that have had a confirmed *Neospora* abortion can be made

with the knowledge that there is a higher risk of repeat abortion in these animals.^{35,52} Seropositive cows also have a greater risk of abortion, and there is a very high probability of congenital infection in their calves.^{35,43,52,57,60} Seropositive heifers also have been shown to have reduced milk production.⁵¹ Epidemiological studies in both dairy and beef cattle have found that seropositive cattle have an increased rate of culling for a variety of reasons.^{50,57}

Although various antimicrobial agents have been tested against *Neospora caninum* in vitro, there is no known method whereby an infected cow can be cleared of the infection.²³

A provisional killed vaccine has recently become available for *Neospora* but there is no information on its efficacy in reducing fetal infection or abortion in an infected cow or in preventing postnatal infection in a non-infected cow.

There are no proven methods available to prevent postnatal infection, as details of the life cycle are still incomplete. However, based on experimental evidence that the dog can be a definitive host³⁴ and the association between dogs on the dairy and seroprevalence in the herd,⁴⁴ it would be prudent to take measures to reduce the potential for this type of transmission. The removal of all potentially infected tissues, such as aborted fetuses and placentas from the environment, that might serve as a source of infection for susceptible hosts would be advisable. In addition, fecal contamination of feed and water sources by other animals should be minimized.

References

1. Abbitt B. Protozoal abortion in cattle. In Current Therapy in Large Animal Theriogenology; RS Youngquest Editor, W. B. Saunders, 1997 pp 383-385.
2. Anderson ML, Blanchard PC, Barr BC, Dubey JP, Hoffman RL, Conrad PA. *Neospora*-like protozoan infection as a major cause of abortion in California dairy cattle. J Am Vet Med Assoc 198;241, 1991
3. Anderson ML, Barr BC, Conrad PA. Protozoal causes of reproductive failure in domestic ruminants. Veterinary Clinics of North America, Food Animal Practice, 1994 Nov, 10(3):439-61.
4. Anderson, ML, Palmer C, Thurmond M, Picanso JP, Blanchard PC, Breitmeyer RE, Layton AW, McAllister M, Daft BM, Kinde H, Read DH, Dubey JP, Conrad PA, Barr BC. Evaluation of abortions in cattle attributable to neosporosis in selected dairy herds in California. J Am Vet Med Assoc, 1995, 207(9):1206-1210.
5. Anderson, ML, Reynolds JP, Rowe JD, Sverlow KW, Packham AE, Barr BC, Conrad PA. Evidence of vertical transmission of *Neospora* infection in dairy cattle. J Am Vet Med Assoc, 210:1169-1172, 1997.
6. Baker DG, Morishita TY, Brooks DL, Shen SK, Lindsay DS, Dubey JP. Experimental oral inoculations in birds to evaluate potential definitive hosts of *Neospora caninum*. J. Parasitol. 81:783-5, 1995.
7. Barr BC, Anderson ML, Dubey JP, Conrad PA. *Neospora*-like protozoal infections associated with bovine abortions. Vet Pathol 28:116, 1991.
8. Barr BC, Conrad PA, Dubey JP, Anderson ML. *Neospora*-like encephalomyelitis in a calf: pathology, ultrastructure, and immunoreactivity. J Vet Diagn Invest 3:39, 1991.
9. Barr BC, Conrad PA, Breitmeyer R, Sverlow K, Anderson ML, Reynolds J, Chauvet AE, Dubey JP, Ardans AA. Congenital *Neospora* infection in calves born from cows that had previously aborted *Neospora*-infected fetuses: Four cases (1990-1992). J Am Vet Med Assoc 202:113, 1993.
10. Barr BC, Rowe JD, Sverlow KW, BonDurant BH, Ardans AA, Oliver MN, Conrad PA. Experimental reproduction of bovine fetal *Neospora* infection and death with a bovine *Neospora* isolate. J Vet Diagn Invest 6:207-215, 1994.
11. Barr BC, Anderson ML, Sverlow KW, Conrad PA. Diagnosis of fetal *Neospora* infection with the *Neospora* indirect fluorescent antibody test (IFAT). Vet Rec, 1995, 137, 611-613.
12. Baszler TV, Knowles DP, Dubey JP, Gay JM, Mathison BA, McElwain TF. Serological diagnosis of bovine neosporosis by *Neospora caninum* monoclonal antibody-based competitive inhibition enzyme-linked immunosorbent assay. Journal of Clinical Microbiology 34:1432-1438, 1996.
13. Bjerkas I, Mohn SF, Presthus J. Unidentified cystforming sporezoan causing encephalomyelitis and myositis in dogs. Z. Parasitenkd 70:271, 1984.
14. Bjerkas I & Dubey JP: Evidence that *Neospora caninum* is identical to the *Toxoplasma*-like parasite of Norwegian dogs. Acta Vet Scand 32:407, 1991.
15. Bjorkman C, Johansson O, Stenlund S, Holmdahl OJM, Uggla A. *Neospora* species infection in a herd of dairy cattle. J AM Vet Med Assoc 208:1441-1444, 1996.
16. Bjorkman C, Holmdahl OJM, Uggla A. An indirect enzyme-linked immunoassay (ELISA) for demonstration of antibodies to *Neospora caninum* in serum and milk of cattle. Veterinary Parasitology 68:251-260, 1997.
17. Bjorkman C, Hemphill A. Characterization of *Neospora caninum* iscom antigens using monoclonal antibodies. Parasitic Immunology 20:73-80, 1998.
18. Campero CM, Anderson ML, Conosciuto G, Odriozola H, Bretschneider G, Poso MA. *Neospora caninum*-associated abortion in a dairy herd in Argentina. Vet Rec 143:228-9, 1998.
19. Conrad PA, Barr BC, Sverlow KW, Anderson ML, Daft BM, Kinde H, Dubey JP, Munson L, Ardans AA. In vitro isolation and characterization of a *Neospora* sp. from aborted bovine fetuses. Parasitology 106:239, 1993.
20. Conrad PA, Sverlow KW, Anderson ML, Rowe J, BonDurant B, Tuter G, Breitmeyer RE, Palmer C, Thurmond M, Ardans AA, Barr BC. Detection of serum antibody responses in cattle with natural or experimental *Neospora* infections. J Vet Diagn Invest 5:572, 1993.
21. Dubey JP, Carpenter JL, Speer CA, Topperrk MJ, Uggla A. Newly recognized fatal protozoan disease of dogs. J Am Vet Med Assoc 192:1269, 1988.
22. Dubey JP, Hattel AL, Lindsay DS, Topper MJ. Neonatal *Neospora caninum* infection in dogs: Isolation of the causative agent and experimental transmission. J Am Vet Med Assoc 193:1259-63, 1988.
23. Dubey JP, Lindsay DS. A review of *Neospora caninum* and neosporosis. Veterinary Parasitology 67:1-59, 1996.
24. Dubey JP, Lindsay DS, Adams DS, Gay JM, Baszler TV, Blagburn BL, Thulliez P. Serologic responses of cattle and other animals infected with *Neospora caninum*. Am J Vet Res 57:329-336, 1996.
25. Fondevila D, Aanor S, Pumarola M, Dubey JP. *Neospora caninum* identification in an aborted fetus in Spain. Veterinary Parasitology 77:187-9, 1998.
26. Gottstein B, Hentrich B, Wyss R, Theur B, Busato A, Steark KD, Meuller N. Molecular and immunodiagnostic investigations on bovine neosporosis in Switzerland. International Journal for Parasitology 28:679-91, 1998.
27. Jenkins MC, Wouda W, Dubey JP. Serological response over time to recombinant *Neospora caninum* antigens in cattle after Neosporosis-induced abortion. Clin and Diagnostic Laboratory Immunology 4:270-274, 1997.
28. Lally NC, Jenkins MC, Dubey JP. Evaluation of two *Neospora caninum* recombinant antigens for use in an enzyme-linked immunosorbent assay for the diagnosis of bovine neosporosis. Clinical and Diagnostic Laboratory Immunology 3:275-279, 1996.
29. Louie K, Sverlow KW, Barr BC, Anderson ML, Conrad PA. Clon-

- ing and characterization of two recombinant *Neospora* protein fragments and their use in serodiagnosis of bovine neosporosis. *Clinical and Diagnostic Laboratory Immunology* 4:692-699, 1997.
30. Lindsay DS, Dubey JP, Cole RA, Nuehring LP, Blagburn BL. *Neospora*-induced protozoal abortions in cattle. *Comp Cont Ed Pract Vet* 15:882, 1993.
 31. Marsh AE, Barr BC, Sverlow KW, Ho M, Dubey JP, Conrad PA. Sequence analysis and comparison of ribosomal DNA from bovine *Neospora* to similar coccidial parasites. *Journal of Parasitology* 81:530-535, 1995.
 32. McAllister MM, Huffman EM, Hietala SK, Conrad PA, Anderson ML, Salman MD. Evidence suggesting a point source exposure in an outbreak of bovine abortion due to neosporosis. *J Vet Diagn Invest* 8:355-357, 1996.
 33. McAllister MM, Jolley WR, Wills RA, Lindsay DS, McGuire AM, Tranas JD. Oral inoculation of cats with tissue cysts of *Neospora caninum*. *Am J Vet Res* 59:441-4, 1998.
 34. McAllister MM, Dubey JP, Lindsay DS, Jolley WR, Wills RA, McGuire AM. Dogs are definitive hosts of *Neospora caninum*. *International Journal for Parasitology* 28:1473-1478, 1998.
 35. Moen AR, Wouda W, Mul MF, Graat EAM, van Werven T. Increased risk of abortion following *Neospora caninum* abortion outbreaks: a retrospective and prospective cohort study in four dairy herds. *Theriogenology* 49:1301-1309, 1998.
 36. Nietfeld JC, Dubey JP, Anderson ML, Libal MC, Yaeger MJ, Neiger RD. *Neospora*-like protozoan infection as a cause of abortion in dairy cattle. *J Vet Diagn Invest* 4:233-226, 1992.
 37. Osawa T, Wastling J, Maley S, Buxton D, Innes EA. A multiple antigen ELISA to detect *Neospora*-specific antibodies in bovine sera, bovine foetal fluids, ovine and caprine sera. *Veterinary Parasitology* 79:19-34, 1998.
 38. Otter A, Jeffrey M, Scholes SFE, Helmick B, Wilesmith JW, Trees AJ. Comparison of histology with maternal and fetal serology for the diagnosis of abortion due to bovine neosporosis. *Vet Rec* 141:487-9, 1997.
 39. Packham AE, Sverlow KW, Conrad PA, Loomis EF, Rowe JD, Anderson ML, Marsh AE, Cray C, Barr BC. A modified agglutination test for *Neospora caninum*: development, optimization, and comparison to the indirect fluorescent-antibody test and enzyme-linked immunosorbent assay. *Clinical and Diagnostic Laboratory Immunology* 5:467-73, 1998.
 40. Paré J, Thurmond MC, Hietala SK. Congenital *Neospora* infection in dairy cattle. *Vet Rec* 134:531-532, 1994.
 41. Paré J, Hietala SK, Thurmond MC. An enzyme-linked immunosorbent assay (ELISA) for serological diagnosis of *Neospora* infection in cattle. *J Vet Diagn Invest* 7:352-359, 1995.
 42. Paré J, Thurmond MC, Hietala SK. Congenital *Neospora caninum* infection in dairy cattle and associated calfhood mortality. *Can J Vet Res* 60:133-9, 1996.
 43. Paré J, Thurmond MC, Hietala SK. *Neospora caninum* antibodies in cows during pregnancy as a predictor of congenital infection and abortion. *J Parasitol.* 83:82-87, 1997.
 44. Paré J, Fecteau G, Fortin M, Marsolais G. Seroepidemiologic study of *Neospora caninum* in dairy herds. *J Am Vet Med Assoc* 213:1595-8, 1998.
 45. Schares G, Peters M, Wurm R, Bearwald A, Conraths FJ. The efficiency of vertical transmission of *Neospora caninum* in dairy cattle analyzed by serologic techniques. *Veterinary Parasitology* 80:87-98, 1998.
 46. Thilsted JP, Dubey JP. Neosporosis-like abortions in a herd of dairy cattle. *J Vet Diagn Invest* 1:205, 1989.
 47. Thornton RN, Thompson EJ, Dubey JP. *Neospora* abortion in New Zealand cattle. *N Z Vet J* 39:129, 1991.
 48. Thornton RN, Gajadhar A, Evans J. *Neospora* abortion epidemic in a dairy herd. *N.Z. Vet J.* 42:190-191, 1994.
 49. Thurmond MC and Hietala SK. Strategies to control *Neospora* infection in cattle. *Bovine Practitioner* 29:60-63, 1995.
 50. Thurmond MC and Hietala SK. Culling associated with *Neospora caninum* infection in dairy cattle. *Am J Vet Res* 57:1559-1562, 1996.
 51. Thurmond MC and Hietala SK. Effect of *Neospora caninum* infection on milk production in first lactation dairy cows. *J Am Vet Med Assoc.* 210:672-674, 1997.
 52. Thurmond MC and Hietala SK. Effect of congenitally acquired *Neospora caninum* infection on risk of abortion and subsequent abortions in dairy cattle. *Am J Vet Res* 58:1381-1385, 1997.
 53. Thurmond MC, Hietala Sk, Blanchard PC. Herd-based diagnosis of *Neospora caninum*-induced endemic and epidemic abortion in cows and evidence for congenital and postnatal transmission. *J Vet Diagn Invest* 9:44-49, 1997.
 54. Thurmond MC, Hietala Sk, Blanchard PC. Predictive values of fetal histopathology and immunoperoxidase staining in diagnosing bovine abortion caused by *Neospora caninum* in a dairy herd. *J Vet Diagn Invest* 11:90-94, 1999.
 55. Thurmond MC and Hietala SK. *Neospora caninum* infection and abortion in cattle. In Howard and Smith eds. *Current Veterinary Therapy: Food animal practice*, 1999, W.B Saunders, pp 425-431.
 56. UgglA, Stenlund OJ, Jakubek EB, Thebo P, Kindahl H, Bjorkman C. Oral *Neospora caninum* inoculation of neonatal calves. *International Journal for Parasitology* 28:1467-72, 1998.
 57. Waldner CL, Jansen ED, Ribble CS. Determination of the association between *Neospora caninum* infection and reproduction performance in beef herds. *J Am Vet Med Assoc* 213:685-90, 1998.
 58. Williams DJL, McGarry J, Guy F, Trees AJ. Novel ELISA for detection of *Neospora*-specific antibodies in cattle. *Vet Rec* 140:328-331, 1997.
 59. Wouda W, Dubey JP, Jenkins MC. Serological diagnosis of bovine fetal neosporosis. *J Parasitol.* 83:545-7, 1997.
 60. Wouda W, Moen AR, Schukken YH. Abortion risk in progeny of cows after a *Neospora caninum* epidemic. *Theriogenology* 49:1311-1316, 1998.
 61. Wouda W, Brinkhof J, van Maanen C, de Gee AL, Moen AR. Serodiagnosis of neosporosis in individual cows and dairy herds: a comparative study of three enzyme-linked immunosorbent assays. *Clinical and Diagnostic Laboratory Immunology* 5:711-6, 1998.