

# Prevention and Prediction of Displaced Abomasum in Dairy Cows

T. Geishauser, K. Leslie, T. Duffield

Department of Population Medicine, Ontario Veterinary College, University of Guelph, Guelph, N1G 2W1, Canada

## Abstract

The objective of this article is to summarize studies that have been conducted on the prevention and prediction of displaced abomasum in dairy cows. Our findings indicate that displaced abomasum is a moderately heritable trait. Subclinical ketosis is a significant risk factor of displaced abomasum. Thus, displaced abomasum incidence might be lowered by genetic selection and by prevention of subclinical ketosis. To predict displaced abomasum, aspartate-aminotransferase activity in blood,  $\beta$ -hydroxybutyrate concentration in blood and milk, and the fat-protein-ratio in milk may be used. All of these parameters are frequently increased prior to the diagnosis of displaced abomasum.

Displaced abomasum (DA) causes economic losses due to treatment costs, discarded milk, decreased milk yield and increased risk of removal from the herd (Martin et al. 1978, Milian-Suazo et al. 1988, Bartlett et al. 1995, Geishauser et al. 1998b). In the lactation in which DA was diagnosed losses ranged from US\$ 250 to US\$ 400 per case, depending on the technique used for surgery (closed or open) (Bartlett et al. 1995). In North America there are about 11 million dairy cows. Given a conservative average DA lactational incidence rate of 2% (Kelton 1995), about 220,000 cases of DA occur per year, resulting in annual losses of about \$55 to 88 M (US). With a higher average incidence rate of 5% (Duffield 1997), about 550,000 cases of DA would occur, causing annual losses of about \$137 to 220 million (US). Considering these economic losses, it is surprising means for prevention and monitoring of DA are not well described (Geishauser 1995).

## Prevention


In a study conducted in Hessia, Germany, DA was most frequently diagnosed in Black Holsteins (Geishauser et al. 1996a). The degree of heritability of DA was estimated at 24% in Black Holstein herds using the similarity between mothers and daughters (Geishauser et al. 1996b). In Ontario, Canada, the degree of heritability of DA was estimated at 28% in Black

Holstein herds, using the similarity between fathers and daughters (Uribe et al. 1995). From these studies, it was suggested that in these regions and breeds, DA is a moderately heritable trait. The incidence of DA may be lowered by genetic selection (Uribe et al. 1995, Geishauser et al. 1996b). It was recommended not to breed cows affected with DA (Geishauser et al. 1996b).


Aspartate-aminotransferase (AST) activity, the concentrations of  $\beta$ -hydroxybutyrate (BHB) and glucose, and body condition score were evaluated three weeks prior to calving, as well as in the first and second week after calving in case or control cows that were subsequently diagnosed with DA or not, respectively. Three weeks prior to calving all of the above parameters were normal in cows diagnosed with DA four to seven weeks later (Fig. 1 to 3). However, AST and BHB were significantly increased in the first and second week after calving in cows diagnosed with DA one to three weeks later as compared to control cows (Fig. 1 and 2) (Geishauser et al. 1998c). If blood AST activity in the first week post partum was  $\geq 1700$  nkat/l, the odds were three to one (compared to  $< 1700$  nkat/l) that DA would be diagnosed one to three weeks later (odds ratio) (in the second week post partum the odds ratio was eight to one). If blood BHB levels were  $\geq 1000$   $\mu\text{mol/l}$ , the odds were two to one (compared to  $< 1000$   $\mu\text{mol/l}$ ) that DA would be diagnosed one to three weeks later (in the second week post partum the odds ratio was four to one) (Geishauser et al. 1997b). Glucose was significantly decreased in the second week after calving, but not in the first week (Fig. 3). Body condition score was unaltered (Geishauser et al. 1998c).

Ketolac<sup>®</sup> BHB (Hoechst, Unterschleißheim/Germany) is a cow-side test that measures  $\beta$ -hydroxybutyrate in milk. This test was evaluated during the first two weeks after calving in cows subsequently diagnosed with DA and in control cows. If Ketolac<sup>®</sup> BHB indicated  $\geq 100$   $\mu\text{mol}$  BHB/l milk in the first two weeks post partum, the odds were three to one (compared to  $< 100$   $\mu\text{mol/l}$ ) that DA would be diagnosed one to three weeks later (Geishauser et al. 1997c).


Daily milk yield, milk fat and protein percentage and milk fat-protein-ratio (FPR) in the first test after calving by the Dairy Herd Improvement (DHI) organi-



**Figure 1.** AST activity in blood three weeks ante partum (a.p.), and in the first and second week post partum (p.p.) in cows subsequently diagnosed with DA (■) and in herdsmates not diagnosed with DA (□). Median and interquartile range are given (Geishauser et al. 1998c).



**Figure 2.** BHB concentration in blood three weeks ante partum (a.p.), and in the first and second week post partum (p.p.) in cows subsequently diagnosed with DA (■) and in herdsmates not diagnosed with DA (□). Median and interquartile range are given (Geishauser et al. 1998c).




**Figure 3.** Glucose concentration in blood three weeks ante partum (a.p.), and in the first and second week post partum (p.p.) in cows subsequently diagnosed with DA (■) and in herdsmates not diagnosed with DA (□). Median and interquartile range are given (Geishauser et al. 1998c).

zation were evaluated in cows subsequently diagnosed with DA and in control cows. One to three weeks prior to DA diagnosis, daily milk yield and milk protein percentage were decreased; milk fat percentage and FPR were increased (Geishauser et al. 1999). If milk FPR was  $\geq 1.4$ , the odds were two to one (compared to  $< 1.4$ ) that DA was diagnosed one to three weeks later (Geishauser et al. 1997a).


BHB concentrations  $\geq 1000 \mu\text{mol/l}$  blood (Työppönen und Kauppinen, 1980), Ketolac<sup>®</sup> BHB findings  $\geq 100 \mu\text{mol}$  BHB/l milk (Geishauser et al. 1998a) and milk FPR  $\geq 1.4$  (Dirksen et al. 1997) may indicate subclinical ketosis. Our findings suggest that subclinical ketosis is a risk factor of DA. Consequently, prevention of subclinical ketosis may prevent DA (Duffield 1997).

### Prediction


As previously discussed, metabolic parameters may be used to predict DA. If AST is used as a test for subsequent DA diagnosis, the true positive rate, and the false positive rate, respectively, depend on the the cut-off level used. The true positive rate gives the sensitivity of the test (rate of positive tests in DA cows); the specificity (rate of negative tests in healthy cows) can be used to calculate the false positive rate ( $1 - \text{specificity}$ ). AST findings  $\geq 1700 \text{ nkat/l}$  in the first week post partum truly indicated subsequent DA diagnosis in 61% (second week post partum 79%) and falsely indicated DA in 46% (second week post partum 31%) (Fig. 4). BHB findings  $\geq 1000 \mu\text{mol/l}$  in the first week post partum truly indicated DA in 61% (second week post partum 64%), and falsely indicated DA in 38% (second week post partum: 31%) (Fig. 5) (Geishauser et al. 1997b). Ketolac<sup>®</sup> BHB findings showing  $\geq 100 \mu\text{mol}$  BHB/l milk truly indicated DA in 46% and falsely indicated DA in 19% (Geishauser et al. 1997c). A FPR  $\geq 1.4$  truly indicated DA in 80% and falsely indicated DA in 31% (Fig. 6) (Geishauser et al. 1997a).



**Figure 4.** AST activity in blood in the first and second week post partum (p.p.) as a test to monitor for DA. True positive rate and false positive rate at different AST cut off levels (Geishauser et al. 1997b).



**Figure 5.** BHB concentration in blood in the first and second week post partum (p.p.) as a test to monitor for DA. True positive rate and false positive rate at different BHB cut off levels (Geishauser et al. 1997b).



**Figure 6.** Fat-protein-ratio in milk in the first test after calving by the Dairy Health Corporation as a test to monitor for DA. True positive rate and false positive rate at different FPR cut off levels (Geishauser et al. 1997a).

Suppose in a herd of 100 dairy cows the lactational incidence rate of DA diagnosis is 2% (Kelton 1995). All cows are monitored for DA on a regular basis in the first month after calving, because 80% of all DA are diagnosed in the first month after calving (Constable et al. 1992). Cows are monitored weekly using BHB in blood. BHB findings  $\geq 1000 \mu\text{mol/l}$  are taken as a positive result. This test truly indicates DA in 60% and falsely indicates DA in 30%. In this herd, two cases of DA will be diagnosed per year. Using blood BHB as a test, one DA will be truly predicted but 30 cases will be falsely predicted (Table 1 and 2). Ketolac findings  $\geq 100 \mu\text{mol BHB/l}$  milk would truly indicate 0.7 DA cases and falsely indicate 19 DA cases. A FPR  $\geq 1.4$  would truly indicate 1.3 DA cases and falsely indicate 30 DA cases. We would like to encourage the reader to perform their own evaluation using a spreadsheet (e.g. Excel 7.0, Microsoft Corporation, Seattle/USA) (Table 1).

Hence, the parameters evaluated here for DA monitoring are insufficiently sensitive (low true positive rate), and insufficiently specific (high false positive rate) under normal conditions. However, under certain conditions monitoring cows for DA weekly in the first month after calving may be profitable when using parameters evaluated (sensitivity 80%, specificity 70%) (Table 3) (Geishauser 1998). These conditions include: DA incidence of 20% (Jacobsen 1995), high losses per case (treatment costs, milk losses, stock losses) (Bartlett et al. 1995), test costs less than \$0.5 US per test, and effective means to prevent DA in cows at risk (administration of glucogenic substances) (Duffield 1997).

Future work may focus on the development of economical tests and effective preventive strategies.

1. Bartlett, P.C., M. Kopcha, P.H. Coe, N.K. Ames, P.L. Ruegg, and R.J. Erskine. 1995. Economic comparison of the pyloro-omentopexy vs the roll-and-toggle procedure for treatment of left displacement of the abomasum in dairy cattle. *J. Am. Vet. Med. Assoc.* 206:1156-1162.
2. Constable, P.D., G.Y. Miller, G.F. Hoffsis, B.L. Hull, and D.M. Rings.

**Table 1.** Structure of a scenario to monitor dairy herds for displaced abomasum (DA) (Geishauser 1998).

A	B	C	D	E	F	G
2				Real	Predicted	
3					True	False
4	Herd Size	100	LDA Cases per Year	$C4 * C5$	$C4 * C5 * C9 * C10$	$C4 * (1 - (C5 * C6)) * C11$
5	LDA Incidence	0,02	LDA Loss per Year	$C4 * C5 * C6$	$F4 * C6$	$G4 * C6$
6	Loss per LDA (\$)	250				
7						
8	Test Period (days)	30	Tests per Year	$C4 * C8 * C12$		
9	LDA% per Test Period	0,8	Test Costs per Year (\$)	$C4 * C8 * C12 * C13$		
10	True Positive Rate	0,6	Prevention (\$) per Year	$C14 * (F4 + G4)$	$C14 * F4$	$C14 * G4$
11	False Positive Rate	0,3	LDA Loss per Year (\$)	$C6 * (E4 - F4)$		
12	Tests per Test Period	0,14	Test+Prevention+Loss (\$)	$E9 + E10 + E11$		
13	Costs per Test (\$)	4				
14	Costs per Prevention (\$)	50	Cost Effect Monitoring	$F5 - E12$		

**Table 2.** Scenario A to monitor dairy herds for displaced abomasum (DA) (Geishauser 1998).

A	B	C	D	E	F	G
2				Real	Predicted	
3					True	False
4	Herd Size	100	LDA Cases per Year	2	1	30
5	LDA Incidence	0,02	LDA Loss per Year	500	240	7380
6	Loss per LDA (\$)	250				
7						
8	Test Period (days)	30	Tests per Year	429		
9	LDA5 per Test Period	0,8	Test Costs per Year (\$)	1714		
10	True Positive Rate	0,6	Prevention (\$) per Year	1524	48	1476
11	False Positive Rate	0,3	LDA Loss per Year (\$)	260		
12	Tests per Test Period	0,14	Test+PreventionLoss (\$)	3498		
13	Costs per Test (\$)	4				
14	Costs per Prevention (\$)	50	Cost Effect Monitoring	-3258		

**Table 3.** Scenario B to monitor dairy herds for displaced abomasum (DA) (Geishauser 1998).

A	B	C	D	E	F	G
2				Real	Predicted	
3					True	False
4	Herd Size	100	LDA Cases per Year	20	13	25
5	LDA Incidence	0,2	LDA Loss per Year	8000	5120	10080
6	Loss per LDA(\$)	400				
7						
8	Test Period (days)	30	Tests per Year	429		
9	LDA% per Test Period	0,8	Test Costs per Year (\$)	214		
10	True Positive Rate	0,8	Prevention (\$) per Year	1900	640	1260
11	False Positive Rate	0,3	LDALoss per Year (\$)		2880	
12	Tests per Test Period	0,14	Test+Prevention+Loss (\$)	4994		
13	Costs per Test (\$)	0,5				
14	Costs per Prevention (\$)	50	Cost Effect Monitoring	126		

1992. Risk factors for abomasal volvulus and left abomasal displacement in cattle. *Am. J. Vet. Res.* 53:1184-1192.

3. Dirksen, G., C. Hagert-Theen, M. Alexander-Katz, and A. Berger. 1997. Stoffwechselüberwachung bei Kühen in der Hochlaktation anhand von Milchparametern. 1. Tagesmilchmenge, Fett- und Eiweißkonzentration, Fett-Eiweiß-Quotient, Harnstoffkonzentration. *Tierärztl. Umsch.* 52:319-324.

4. Duffield, T. 1997. Effects of a monensin controlled release capsule on energy metabolism, health and production in lactating dairy cattle. Guelph, University, DVSc-Thesis.

5. Geishauser, T. 1995. Untersuchungen zur Labmagenmotorik von Kühen mit Labmagenverlagerung. Verlag Shaker, Aachen, ISBN 3-8265-0757-6.

6. Geishauser, T. 1998. Vorbeuge und Früherkennung von Labmagenverlagerung bei Milchkuhen. *Tierärztl. Umsch.* 53:601-606.

7. Geishauser, T., M. Diederichs, and K. Failing. 1996a. Vorkommen von Labmagenverlagerung bei Rindern in Hessen. *Dtsch. Tierärztl. Wschr.* 103:142-144.

8. Geishauser, T., M. Diederichs, and R. Beuing. 1996b. Schätzung der Erbllichkeit von Labmagenverlagerung bei Deutsch-Schwarz-bunten Rindern in Hessen. *Zbl. Vet. Med. A* 43, 87-92.

9. Geishauser, T., K. Leslie, T. Duffield, and V. Edge 1997a. Fat/protein ratio in first DHI test milk as tests for displaced abomasum in dairy cows. *Zbl. Vet. Med. A* 44:265-270.

10. Geishauser, T., K. Leslie, T. Duffield, and V. Edge 1997b. An evaluation of aspartate-aminotransferase activity and  $\beta$ -hydroxybutyrate concentration in blood as tests for left displaced abomasum in dairy cows. *Am. J. Vet. Res.* 58:1216-1220.

11. Geishauser, T., K. Leslie, T. Duffield, and V. Edge. 1997c. An evaluation of milk ketone tests for prediction of left displaced abomasum in dairy cows. *J. Dairy Sci.* 80:3188-3192.

12. Geishauser, T., K. Leslie, D. Kelton, and V. Edge. 1998a. An evaluation of five cow-side tests for use with milk to detect subclinical ketosis in dairy cows. *J. Dairy Sci.* 81:438-443.

13. Geishauser, T., M. Shoukri, D. Kelton, and K. Leslie 1998b. Analysis of survivorship after left displaced abomasum is diagnosed in dairy cows. *J. Dairy Sci.* 81:2346-2353.

14. Geishauser, T., K. Leslie, T. Duffield, D. Sandals, and V. Edge. 1998c. The association between selected metabolic parameters and left abomasal displacement in dairy cows. *Zbl. Vet. Med. A* 45:499-511.

15. Geishauser, T., K. Leslie, T. Duffield, and V. Edge. 1999. The as-

sociation between first DHI test milk parameters and displaced abomasum in dairy cows. *Berl. Münch. Tierärztl. Wschr.* 112: in press.

16. Jacobsen, K.L. 1995. Displaced abomasa and thin cows in a component-fed dairy herd. *Food Animal Medicine and Management* 17:8, 21-27.

17. Kelton, D.F. 1995. Monitoring, and investigating the relationships among, health, management, productivity and profitability on Ontario dairy farms. Guelph, University, PhD-Thesis.

18. Martin, W., K.L. Kirby, and R.A. Curtis. 1978. Left abomasal displacement in dairy cows. Its relationship to production. *Can. Vet. J.*

19:250-253.

19. Milian-Suazo, F., H. N. Erb, and R. D. Smith. 1988. Descriptive epidemiology of culling in cows from 34 herds in New York state. *Prev. Vet. Med.* 6:243-251.

20. Tyoppönen, J., and K. Kauppinen. 1980. The stability and automatic determination of ketone bodies in blood samples taken in field conditions. *Acta Vet. Scand.* 21:55-61.

21. Uribe, H.A., B.W. Kennedy, S.W. Martin, and D.F. Kelton. 1995. Genetic parameters for common health disorders of Holstein cows. *J. Dairy Sci.* 78:421-430.