after vaccination with cytopathic bovine viral diarrhea virus. Am J Vet Res 49:1040-1044,1988. 4. Bolin SR, Littledike ET, Ridpath JF: Serologic detection and practical consequences of antigen diversity among bovine viral diarrhea viruses in a vaccinated herd. Am J Vet Res 52:1033-1037, 1991. 5. Bolin SR, Moennig V, Gourley NE, et al: Monoclonal antibodies with neutralizing activity segregate isolates of bovine viral diarrhea virus into groups. Arch Virol 99:L117-123, 1988. 6. Bolin SR, Ridpath JF: Range of viral neutralizing activity and molecular specificity of antibodies induced in cattle by inactivated bovine viral diarrhea virus vaccines. Am J Vet Res 51:703-707, 1990. 7. Bolin SR, Ridpath JF: Specificity of neutralizing and precipitating antibodies induced in healthy calves by monovalent modified-live bovine viral diarrhea virus vaccines. Am J Vet Res 50:817-821, 1989. 8.Bolin SR, McClurkin AW, Cutlip RC, et al: Severe clinical disease induced in cattle persistently infected with noncytopathic bovine viral diarrhea virus by superinfection with cytopathic bovine viral diarrhea virus. Am J Vet Res 46:573-578, 1985. 9.Bolanger D, Waxweiler S, Karelle L, et al: Characterization of monoclonal antibodies to bovine viral diarrhoea virus: Evidence of a neutralizing activity against gp48 in the presence of goat anti-mouse J Gen Virol 72:1195-1198, immunoglobulin serum. 1991. 10. Castrucci G, Avellini G, Cilli V, et al: A study of immunologic relationships among serologically heterologous strains of bovine viral diarrhea virus by cross immunity tests. Cornell Vet 65:65-72, 1975. 11. Collett MC: Molecular genetics of pestiviruses. Comp Immunol Infect Dis 15:145-154, 1992. 12. Deregt D, Masri SA, Cho HJ, et al: Monoclonal antibodies to the p80/125 and gp53 proteins of bovine viral diarrhea virus: their potential use as diagnostic reagents. Can J Vet Res 54:343-348, 1990. 13. Donis RO, Corapi W, Dubovi EJ: Neutralizing monoclonal antibodies to bovine viral diarrhea virus bind to the 56K to 58K glycoprotein. J Gen Virol 69:77-86, 1988. 14.Edwards S, Wood L, Hewitt-Taylor C, et al: Evidence for an immunocompromising effect of bovine pestivirus on bovid herpevirus 1 vaccination. Vet Res Comm 10:297-302, 1986. 15. Gunn HM: Role of fomites and flies in the transmission of bovine viral diarrhoea virus. Vet Rec 132:548-585, 1993. 16. Howard CJ, Clarke MC, Sopp P, et al: Immunity to bovine virus diarrhoea virus in calves: the role of different T-cell subpopulations analysed by specific depletion in vivo with monoclonal antibodies. Vet Immunol Immunopathol 32:303-314, 1992. 17. Kamstrap S, Roensholt L, Dalsgaard K: Immunological

reactivity of bovine viral diarrhoea virus proteins after proteolytic treatment. Arch Virol S3:225-230, 1991. 18. Kwang J, Littledike ET, Donis RO, et al: Recombinant polypeptide from the gp48 region of bovine viral diarrhea virus (BVDV) detects serum antibodies in vaccinated and infected cattle. Vet Microbiol 32:281-292, 1992. 19. Martin S, Berlinski P, Guimond P, et al: Evaluation of various BVDV isolates to induce protection against challenge with virulent BVDV 890. In Proceedings of the Conference Res Workers Anim Dis 75:183, 1994. 20. McClurkin AW, Littledike ET, Cutlip RC, et al: Production of cattle immunotolerant to bovine viral diarrhea virus. Can J Comp Md 48:158-161, 1984. 21. Nettleman PF, Entrican, G: Ruminant Pestiviruses. Br Vet J. 151:615-642. 1995. 22.Paton DJ, Ibata G, Edwards S, Et al: An ELISA detecting antibody to conserved pestivirus epitopes. J Virol Methods 31:315-324, 1991. 23. Pellerin C, Hurk JVD, Lecomte J, et al: Identification of a new group of bovine viral diarrhea virus strains associated with severe outbreaks and high mortalities. Virology 203:260-268, 1994. 24. Potgieter LND, McCracken MD, Hopkins FM, et al: Effect of bovine viral diarrhea virus infection on the distribution of infectious bovine rhinotracheitis virus in calves. Am J Vet Res 45:687-690, 1984a. 25. Potgeiter LND, McCracken MD, Hopkins FM, et al: Experimental production of respiratory tract disease with bovine viral diarrhea virus. Am J Vet Res 45:1582-1585, 1984b. 26. Potgieter LND, McCracken MD, Hopkins FM, et al: Comparison of the pneumopathogenicity of two strains of bovine viral diarrhea virus. Am J Vet Res 46-151-153, 1985. 27. Potgeiter LND: Immunology of bovine viral diarrhea virus. In: Veterinary Clinics of North America: Food Animal Practice. 11:501-520, 1995. 28. Rumenapf T, Stark R, Meyers G, et al: Structural proteins of hog cholera virus expressed by vaccinia virus: Further characterization and induction of protective immunity. J Virol 65:589-597, 1991. 29. Steck F. Lazarv S. Fev H. et al. Immune responsiveness in cattle fatally affected by bovine virus diarrhea-mucosal disease. Zentrabl for Veterinaermed B27:429-445, 1980. 30. Van Oirschot JT. Congenital infections with non-arbo togaviruses. Vet Microbiol. 8:321-361, 1983. 31. Wensvoort G, Terpstra C, de Kluyver EP: Characterization of porcine and some ruminant pestiviruses by cross-neutralization. Vet Microbiol 20:291-306, 1989. 32. Xue W, Blecha F, Minocha HC: Antigenic variations in bovine viral diarrhea viruses detected by monoclonal antibodies. J Clin Microbiol 28:1688-1693, 1990.

## Abstract

## Preferential sites for arterial blood sampling in cattle

## S. Muylle, P. Anthone, P. Simoens, H. Lauwers

Veterinary Record (1996); 139, 86-88

The aim of this study was to compare the feasibility and reliability of different methods for obtaining samples of arterial blood from calves and adult dairy cows. The intermediate branch of the caudal auricular artery was easy to use in animals of all ages. The common carotid artery was also suitable, but only in standing animals. In recumbent animals, the brachial artery and the common palmar digital artery were suitable in calves up to six months old, and the saphenous artery was suitable in both young and adult cattle. The facial, axillary and median caudal arteries were either difficult to use or unreliable.