

The what, why, and physiologic cost of leaky gut syndrome

S. Rodriguez-Jimenez, DVM, MS; E. A. Horst; E. J. Mayorga, MS; S. K. Kvidera, PhD; M. A. Abeyta; B. M. Goetz; S. Carta, MS; L. H. Baumgard, MS, PhD

Iowa State University, Department of Animal Science, Ames, IA 50011

Corresponding author: Dr. L. H. Baumgard, 313 Kildee Hall, Ames, IA, Email: baumgard@iastate.edu

Abstract

Heat stress and metabolic maladaptation to lactation (ketosis) are 2 economically devastating hurdles to profitability. These stressors affect herds of all sizes and almost every dairy region in the world. The biology of heat stress and ketosis has been studied for almost a half century, but the negative impacts of both are as evident today as they were 30 years ago. Our recent discoveries suggest that endotoxin is the common culprit in both disorders and the intestine appears to be the etiological origin of both metabolic disorders. Endotoxin stimulates the immune system and activated leukocytes switch their metabolism away from oxidative phosphorylation to rely more on aerobic glycolysis. In multiple species, we estimate that immune activation consumes about 0.45 g glucose/lb (1 g/kg) BW^{0.75} or about 0.9 lb (2 kg) glucose/day in an adult lactating dairy cow. Thus, an activated immune system reprioritizes nutrient partitioning away from the synthesis of economically valuable products.

Key words: cow, dairy, leaky gut, metabolism, LPS

Résumé

Le stress dû à la chaleur et les troubles métaboliques associés à la lactation (acidose) représentent deux écueils économiquement désastreux à la profitabilité. Ces deux sources de stress affectent les troupeaux de toutes tailles dans presque toutes les régions laitières du monde. Les fondements biologiques du stress dû à la chaleur et de l'acidose ont été étudiés depuis plus d'un demi-siècle mais leurs impacts négatifs sont aussi évidents aujourd'hui qu'ils ne l'étaient il y a 30 ans. Nos découvertes récentes suggèrent qu'une endotoxine est un facteur commun aux deux conditions et l'intestin semble être l'origine étiologique des deux troubles métaboliques. L'endotoxine stimule le système immunitaire et les leucocytes activés s'éloignent de leur métabolisme basé sur la phosphorylation oxydative et se tournent davantage vers la glycolyse aérobie. Pour plusieurs espèces, nous estimons que l'activation immunitaire consomme à peu près 0.45 g glucose/lb (1g/kg) PC^{0.75} ou près de 0.9 lb (2 kg) de glucose par jour chez une vache laitière adulte en lactation. Ainsi, l'activation du système immunitaire redistribue la division des nutriments et l'éloigne de la synthèse de produits économiquement valables.

Introduction

Suboptimal milk yield limits the USA dairy industry's productive competitiveness, marginalizes efforts to reduce inputs into food production, and increases animal agriculture's carbon footprint. There are a variety of circumstances in a cow's life which result in hindered productivity including heat stress, ketosis, rumen and hindgut acidosis, feed restriction, and psychological stress associated with normal animal practices (i.e., pen changes, weaning, shipping). Although these insults have different origins, a commonality among them is increased production of inflammatory biomarkers and markedly altered nutrient partitioning. We and others have generated preliminary data strongly implicating intestinally derived lipopolysaccharide (LPS) as a culprit in these situations.

Heat Stress

Heat stress (HS) affects blood flow which is diverted from the viscera to the periphery in an attempt to dissipate heat leading to intestinal hypoxia.²⁶ Enterocytes are particularly sensitive to hypoxia and nutrient restriction,⁷⁰ resulting in ATP depletion and increased oxidative and nitrosative stress.²⁷ This contributes to tight junction dysfunction and gross morphological changes that ultimately reduce intestinal barrier function,^{47,66} resulting in increased passage of luminal content into portal and systemic blood.^{27,66} Endotoxin, otherwise referred to as LPS, is a glycolipid embedded in the outer membrane of gram-negative bacteria, which are abundant and prolific in luminal content, and is a well-characterized potent immune stimulator in multiple species.^{5,19,78} Immune system activation occurs when LPS binding protein (LBP) initially binds LPS and together with CD14 and TLR4 delivers LPS for removal and detoxification, thus LBP is frequently used as a biomarker for LPS infiltration.¹² For a detailed description of how livestock and other species detoxify LPS see our recent review.⁵⁶ Endotoxin infiltration into the bloodstream during HS, which was first observed by Graber et al,²² is common among heat stroke patients⁴⁹ and is thought to play a central role in heat stroke pathophysiology as survival increases when intestinal bacterial load is reduced or when plasma LPS is neutralized.^{9,18} It is remarkable how animals suffering from heat stroke or severe endotoxemia share many physiological and metabolic similarities to HS, such as an

increase in circulating insulin.⁵³ Intramammary LPS infusion increased (~2 fold) circulating insulin in lactating cows.⁸² In addition, we intravenously infused LPS into growing calves and pigs and demonstrated >10-fold increase in circulating insulin.^{42,45,68} Interestingly, increased insulin occurs prior to increased inflammation and the temporal pattern agrees with our previous *in vivo* data and a recent *in vitro* report⁷ suggesting LPS stimulates insulin secretion, either directly or via GLP-1.³⁶ The possibility that LPS increases insulin secretion likely explains the hyperinsulinemia we have repeatedly reported in a variety of HS agriculture models.⁴ Again, the increase in insulin in both models is energetically difficult to explain as feed intake was severely depressed in both experiments.

Ketosis and the Transition Period

Recently, the concept that LPS impacts normal nutrient partitioning and potentially contributes to metabolic maladaptation to lactation has started to receive attention. Although LPS itself has not been the primary causative focus, general inflammation has been the topic of investigations. Increased inflammatory markers following parturition have been reported in cows.^{2,6,32,61} Presumably, the inflammatory state following calving disrupts normal nutrient partitioning and is detrimental to productivity,^{6,54} and this assumption was recently reinforced when TNF α infusion decreased productivity (albeit without overt changes in metabolism).^{58,85}

Additionally, in late-lactation cows, injecting TNF α increased (>100%) liver TAG content without a change in circulating NEFA.⁸ Our recent data demonstrates increased inflammatory markers in cows diagnosed with ketosis only and no other health disorders. In comparison with healthy controls, ketotic cows had increased circulating LPS prior to calving and postpartum acute phase proteins such as LBP, serum amyloid A, and haptoglobin were also increased¹ (Figure 1). Endotoxin can originate from a variety of locations, and obvious sources in transitioning dairy cows include the uterus (metritis) and mammary gland (mastitis).⁵⁶ Additionally, we believe intestinal hyperpermeability may also be responsible for periparturient inflammation in dairy cows as many of the characteristic responses (rumen acidosis, decreased feed intake, and psychological stress) occurring during this time can compromise gut barrier function.

Rumen and Hindgut Acidosis

A transitioning dairy cow undergoes a post-calving diet shift from a mainly forage-based to a high-concentrate ration. This has the potential to induce rumen acidosis (RA) as increases in fermentable carbohydrates and DMI stimulate the buildup of short-chain fatty acids and lactic acid.^{16,62} Rumen acidosis has direct and ancillary consequences accompanied by various production issues (decreased DMI, reduced milk yield, milk fat depression) and health challenges such as lami-




Figure 1. Markers of inflammation in healthy (solid line) and ketotic (dashed line) transition cows.

nititis, liver abscesses, and potentially death.^{40,62} The mechanisms linking RA and the development of health disorders are not entirely clear; however, recent literature has indicated that inflammation associated with epithelial damage and consequential LPS translocation are at least partially responsible for production losses associated with RA.^{21,7} Although many hypothesize LPS translocation occurs at the rumen epithelium directly,^{25,60} others point towards LPS translocation in the hindgut to be a potential source of peripheral inflammation.⁵¹ Interestingly, when RA was induced using either alfalfa pellets or high-grain diets, increased peripheral inflammation was only observed in the high-grain group, irrespective of rumen acidotic conditions being similar between the 2 treatments.^{37,38} It was hypothesized that the grain-supplemented group likely had increased starch flow to the hindgut, and therefore, increased fermentation that could potentially lead to hindgut acidosis and LPS translocation across the large intestine. However, we were unable to recreate production losses and systemic inflammation when we abomasally infused 500 g/d of resistant starch⁶⁴ or even 8.8 lb (4 kg)/d of purified corn starch (Abeyta and Baumgard, unpublished). Both of our aforementioned experiments were accompanied with marked reductions in fecal pH, so it is unlikely that large intestinal acidosis per se is the specific reason for systemic inflammation described in the previous reports.^{37,38,51}

Feed Restriction and Psychological Stress

Stress associated with feed restriction along with several other regular production practices (e.g., heat stress, weaning, transportation, overcrowding, restraint, social isolation/mixing) is frequently encountered in animal agriculture¹³ and is associated with gastrointestinal permeability. In fact, we have repeatedly reported reduced intestinal barrier integrity in pigs pair-fed to their HS counterparts.^{66,72} Furthermore, we recently demonstrated shortened ileum villous height and crypt depth, indicating reduced intestinal health, in cows fed 40% of ad libitum intake.⁴⁶ Recent literature indicates that the corticotropin releasing factor (CRF) system may be the mechanism involved in stress-induced leaky gut.^{79,83} The CRF and other members of the CRF signaling family, including urocortin (1, 2, and 3) and their G-protein couple receptors CRF₁ and CRF₂, have been identified as the main mediators of the stress-induced intestinal changes including inflammation, altered intestinal motility and permeability, as well as shifts in ion, water, and mucus secretion and absorption (as reviewed by Rodiño-Janeiro et al⁶⁹). These alterations appeared to be regulated in large part by intestinal mast cells.⁷¹ Mast cells are important mediators of both innate and adaptive immunity and express receptors for the neuropeptides CRF1 and CRF2, which may in part explain the association between emotional stress and intestinal dysfunction.^{3,74} Furthermore, mast cells synthesize a variety of pro-inflammatory mediators (i.e., IFN- γ and TNF- α) that are released upon activation, mainly via degranulation.¹⁴ Excessive mast cell degranulation plays

an important role in the pathogenesis of different intestinal inflammatory disorders.^{71,74} A better understanding of the role psychosocial stress plays on the initiation of different intestinal disorders in livestock is of great interest for animal agriculture systems.

Metabolism of Inflammation

Inflammation induced by LPS has an energetic cost that redirects nutrients away from anabolic processes that support milk and muscle synthesis (see review by Johnson^{34,35}) and thus compromises productivity. Upon activation, immune cells become obligate glucose utilizers via a metabolic shift from oxidative phosphorylation to aerobic glycolysis, a process known as the Warburg effect (Figure 2). This metabolic shift allows for rapid ATP production and synthesis of important intermediates which support proliferation and production of reactive oxygen species.^{10,65} In an effort to facilitate glucose uptake, immune cells become more insulin sensitive and increase expression of GLUT3 and GLUT4 transporters,^{57,63} whereas peripheral tissues become insulin resistant.^{52,67} Furthermore, metabolic adjustments including hyperglycemia or hypoglycemia (depending upon the stage and severity of infection) increased circulating insulin and glucagon, skeletal muscle catabolism and, subsequent nitrogen loss (Figure 3; Wannemacher et al⁸⁴), and hypertriglyceridemia^{17,48,59,84} occur. Interestingly, despite hypertriglyceridemia, circulating BHB often decreases following LPS administration.^{23,43,80,81} The mechanism of LPS-induced decreases in BHB has not been fully elucidated but may be explained by increased ketone oxidation by peripheral tissues.⁸⁶ In addition to changes in circulating metabolites, LPS has been shown to increase liver lipid accumulation, both directly through changes in lipid oxidation and transport enzymes, and indirectly through increases in circulating NEFA.⁸ Collectively, these metabolic alterations are presumably employed to ensure adequate glucose delivery to activated leukocytes.

Energetic Cost of Immune Activation

Immunoactivation leads substantial energetic costs, but the ubiquitous nature of the immune system makes quantifying the energetic demand difficult. Our group recently employed a series of LPS-euglycemic clamps to quantify the energetic cost of an activated immune system. Using this model, we estimated approximately 2.2 lb (1 kg) of glucose is used by an intensely activated immune system during a 12-hour period in lactating dairy cows. Interestingly, on a metabolic body weight basis the amount of glucose utilized by LPS-activated immune system in mid- and late-lactation cows, growing steers and growing pigs were 0.64, 1.0, 1.0, and 1.1 g glucose/kg (2.2 lb) BW^{0.75}/h, respectively.^{28,42,44,45} A limitation to our model is the inability to account for liver's contribution to the circulating glucose pool (i.e., glycogenolysis and gluconeogenesis). However, both glycogenolytic




Figure 2. Metabolic pathway of a resting (A) vs activated (B) leukocyte.




Figure 3. LPS induced alterations in peripheral metabolism.

and gluconeogenic rates have been shown to be increased during infection.^{75,80} Furthermore, we have observed both increased circulating glucagon and cortisol (indirect markers of hepatic glucose output) following LPS administration^{29,30} suggesting we are underestimating the energetic cost of immunoactivation. The reprioritization of glucose trafficking during immunoactivation has particular consequences during lactation, as it requires ~72 g of glucose for synthesizing 2.2 lb (1 kg) milk.

Increased immune system glucose utilization occurs simultaneously with infection-induced decreased feed intake: this coupling of enhanced nutrient requirements with hypophagia obviously decreases the amount of nutrients

available for the synthesis of valuable products (milk, meat, fetus, wool). We and others have now demonstrated that HS, rumen acidosis, and psychological stress increase circulating markers of endotoxin and inflammation. We believe that the circulating LPS originates from the intestine and initiates an immune response. This activated systemic immune response reprioritizes the hierarchy of glucose utilization, and milk synthesis is consequently deemphasized.

Conclusion

In a cow's life there are various situations that hinder animal performance (i.e., heat stress, feed restriction, rumen

acidosis, etc.) and we suggest, based upon the literature and on our supporting evidence, that LPS (of intestinal origin) may be the common culprit in these circumstances. Immune activation in response to LPS markedly alters nutrient partitioning as a means of fueling the immune response. More research is still needed to understand the mechanisms and consequences of intestinal permeability and associated inflammation in order to provide foundational information for developing strategies aimed at maintaining productivity. Furthermore, it is of interest to further elucidate the contribution of inflammation to subclinical hypocalcemia frequently observed in postpartum cows.

Acknowledgement

Parts of this manuscript were first published in the Proceedings of the 2016, 2017, and 2018 Southwest Nutrition Conference in Tempe, AZ.

References

1. Abuajamieh M, Kvidera SK, Fernandez MV, Nayeri A, Upah NC, Nolan EA, Lei SM, DeFrain JM, Green HB, Schoenberg KM, Trout EB, Baumgard LH. Inflammatory biomarkers are associated with ketosis in periparturient Holstein cows. *Res Vet Sci* 2016; 109:81-85.
2. Ametaj BN, Bradford BJ, Bobe G, Nafikov RA, Lu Y, Young JW, Beitz DC. Strong relationships between mediators of the acute phase response and fatty liver in dairy cows. *Can J Anim Sci* 2005; 85:165-175.
3. Ayyadurai S, Gibson AJ, D'Costa S, Overman EL, Sommerville LJ, Poopal AC, Mackey E, Li Y, Moeser AJ. Corticotropin-releasing factor receptor subtype 1 is a critical modulator of mast cell degradation and stress-induced pathophysiology. *J Leukoc Biol* 2017; 102:1299-1312.
4. Baumgard LH, Rhoads RP. Effects of heat stress on postabsorptive metabolism and energetics. *Annu Rev Anim Biosci* 2013; 1:311-337.
5. Berczi I, Bertok L, Bereznai T. Comparative studies on the toxicity of *Escherichia coli* lipopolysaccharide endotoxin in various animal species. *Can J of Microbiol* 1966; 12:1070-1071.
6. Bertoni G, Trevisi E, Han X, Bionaz M. Effects of inflammatory conditions on liver activity in puerperium period and consequences for performance in dairy cows. *J Dairy Sci* 2008; 91:3300-3310.
7. Bhat UG, Ilievski V, Unterman TG, Watanabe K. *Porphyromonas gingivalis* lipopolysaccharide (LPS) upregulates insulin secretion from pancreatic beta cells line MIN6. *J Periodontol* 2014; 85:1629-1636.
8. Bradford BJ, Mamedova LK, Minton JE, Drouillard JS, Johnson BJ. Daily injection of tumor necrosis factor- α increases hepatic triglycerides and alters transcript abundance of metabolic genes in lactating dairy cattle. *J Nutr* 2009; 139:1451-1456.
9. Bynum G, Brown J, Dubose D, Marsili M, Leav I, Pistole TG, Hamlet M, LeMaire M, and Caleb B. Increased survival in experimental dog heatstroke after reduction of gut flora. *Aviat Space Environ Med* 1979; 50:816-819.
10. Calder PC, Dimitriadis G, Newsholme P. Glucose metabolism in lymphoid and inflammatory cells and tissues. *Curr Opin Clin Nutr Metab Care* 2007; 10:531-540.
11. Carlstedt F, Eriksson M, Kiiski R, Larsson A, Lind L. Hypocalcemia during porcine endotoxemic shock: Effects of calcium administration. *Crit Care Med* 2000; 28:2909-2914.
12. Ceciliani F, Ceron JJ, Eckersall PD, Sauerwein H. Acute phase proteins in ruminants. *J Proteomics* 2012; 75:4207-4231.
13. Chen Y, Arsenault R, Napper S, Griebel P. Models and methods to investigate acute stress responses in cattle. *Animals (Basel)* 2015; 5:1268-1295.
14. de Punder K, Pruijboom L. Stress induces endotoxemia and low-grade inflammation by increasing barrier permeability. *Front Immunol* 2015; 6:223.
15. Eckel EF, Ametaj BN. Invited review: Role of bacterial endotoxins in the etiopathogenesis of periparturient diseases of transition dairy cows. *J Dairy Sci* 2016; 99:5967-5990.
16. Enemark JMD. The monitoring, prevention and treatment of sub-acute ruminal acidosis (SARA): A review. *Vet J* 2008; 176:32-43.
17. Filkins JP. Phases of glucose dyshomeostasis in endotoxemia. *Circ Shock* 1978; 5:347-355.
18. Gathiram P, Wells MT, Brock-Utne JG, Gaffin SL. Antilipopolysaccharide improves survival in primates subjected to heat stroke. *Circ Shock* 1987; 23:157-164.
19. Giri SN, Emau P, Cullor JS, Stabenfeldt GH, Bruss ML, Bondurant RH, Osburn BI. Effects of endotoxin infusion on circulating levels of eicosanoids, progesterone, cortisol, glucose and lactic acid, and abortion in pregnant cows. *Vet Microbiol* 1990; 21:211-231.
20. Goff JP, Liesegang A, Horst RL. Diet-induced pseudohypoparathyroidism: A hypocalcemia and milk fever risk factor. *J Dairy Sci* 2014; 97:1520-1528.
21. Gozho, GN, Plaizier JC, Krause DO, Kennedy AD, Wittenberg KM. Subacute Ruminal Acidosis Induces Ruminal Lipopolysaccharide Endotoxin Release and Triggers an Inflammatory Response. *J Dairy Sci* 2005; 88:1399-1403.
22. Graber CD, Reinhold RB, Breman JG, Harley RA, Hennigar GR. Fatal heat stroke. Circulating endotoxin and gram-negative sepsis as complications. *JAMA* 1971; 216:1195-1196.
23. Graugnard DE, Moyes KM, Trevisi E, Khan MJ, Keisler D, Drackley JK, Bertoni G, Looor JJ. Liver lipid content and inflammometabolic indices in periparturient dairy cows are altered in response to prepartal energy intake and postpartal intramammary inflammatory challenge. *J Dairy Sci* 2013; 96:918-935.
24. Griel LC, Zarkower A, Eberhart RJ. Clinical and clinico-pathological effects of *Escherichia coli* endotoxin in mature cattle. *Can J Comp Med* 1975; 39:1-6.
25. Guo J, Chang G, Zhang K, Xu L, Jin D, Bilal MS, Shen X. Ruminal-derived lipopolysaccharide provoked inflammatory injury in the liver of dairy cows fed a high-concentrate diet. *Oncotarget* 2017; 8:46769-46780.
26. Hall DM, Baumgardner KR, Oberley TD, Gisolfi CV. Splanchnic tissues undergo hypoxic stress during whole body hyperthermia. *Am J Physiol* 1999; 276: G1195-G1203.
27. Hall DM, Buettner GR, Oberley LW, Xu L, Matthes RD, Gisolfi CV. Mechanisms of circulatory and intestinal barrier dysfunction during whole body hyperthermia. *Am J Physiol Heart Circ Physiol* 2001; 280:H509- H521.
28. Horst EA, Kvidera SK, Mayorga EJ, Shouse CS, Al-Qaisi M, Dickson MJ, Ydstie J, Ramirez Ramirez HA, Keating AF, Dickson DJ, Griswold KE, Baumgard LH. Effect of chromium on bioenergetics and leukocyte dynamics following immunoactivation in lactating Holstein cows. *J Dairy Sci* 2018; 101:5515-5530.
29. Horst EA, Mayorga EJ, Portner SL, Al-Qaisi M, McCarthy CS, Abeyta MA, Goetz BM, Ramirez-Ramirez HA, Kleinschmit DH, Baumgard LH. Effects of dietary zinc on energetic requirements of an activated immune system following lipopolysaccharide challenge in lactating cows. *J Dairy Sci* 2018; 101 (Suppl. 2): 271.
30. Horst EA, Mayorga EJ, Portner SL, Al-Qaisi M, McCarthy CS, Abeyta MA, Goetz BM, Ramirez-Ramirez HA, Kleinschmit DH, Baumgard LH. Effects of dietary zinc source on inflammatory biomarkers and PMN function following lipopolysaccharide challenge in lactating cows. *J Dairy Sci* 2018; (Suppl. 2): 383.
31. Horst EA, Mayorga EJ, Al-Qaisi M, Abeyta MA, Portner SL, McCarthy CS, Goetz BM, Ramirez-Ramirez HA, Baumgard LH. Effects of maintaining eucalcemia following immunoactivation in lactating cows. *J Dairy Sci* 2018; (Suppl. 2):383.
32. Humblet MF, Guyot H, Boudry B, Mbayahi F, Hanzen C, Rollin F, Godeau JM. Relationship between haptoglobin, serum amyloid A, and clinical status in a survey of dairy herds during a 6-month period. *Vet Clin Pathol* 2006; 35:188-193.
33. Jing L, Zhang R, Liu Y, Zhu W, Mao S. Intravenous lipopolysaccharide challenge alters ruminal bacterial microbiota and disrupts ruminal metabolism in dairy cattle. *Br J Nutr* 2014; 112:170-182.
34. Johnson RW. Inhibition of growth by pro-inflammatory cytokines: An integrated view. *J Anim Sci* 1997; 75:1244-1255.
35. Johnson RW. Immune and endocrine regulation of food intake in sick animals. *Dome Animal Endo* 1998; 15:309-319.

36. Kahles F, Meyer C, Möllmann J, Diebold S, Findeisen HM, Leberherz C, Trautwein C, Koch A, Tacke F, Marx N, Lehrke M. GLP-1 Secretion Is Increased by Inflammatory Stimuli in an IL-6-Dependent Manner, Leading to Hyperinsulinemia and Blood Glucose Lowering. *Diabetes* 2014; 63:3221-3229.
37. Khafipour E, Krause DO, Plaizier JC. A grain-based subacute ruminal acidosis challenge causes translocation of lipopolysaccharide and triggers inflammation. *J Dairy Sci* 2009; 92:1060-1070.
38. Khafipour E, Krause DO, Plaizier JC. Alfalfa pellet-induced subacute ruminal acidosis in dairy cows increases bacterial endotoxin in the rumen without causing inflammation. *J Dairy Sci* 2009; 92:1712-1724.
39. Kimura K, Reinhardt TA, Goff JP. Parturition and hypocalcemia blunts calcium signals in immune cells of dairy cattle. *J Dairy Sci* 2006; 89:2588-2595.
40. Kleen JL, Hooijer GA, Rehage J, Noordhuizen JPTM. Subacute ruminal acidosis (SARA): A review. *J Vet Med* 2003; 50:406-414.
41. Kronfeld DS. Major metabolic determinants of milk volume, mammary efficiency, and spontaneous ketosis in dairy cows. *J Dairy Sci* 1982; 65:2204-2212.
42. Kvidera SK, Horst EA, Abuajamieh M, Mayorga EJ, Sanz-Fernandez MV, Baumgard LH. Technical note: A procedure to estimate glucose requirements of an activated immune system in steers. *J Anim Sci* 2016; 94:4591-4599.
43. Kvidera SK, Dickson MJ, Abuajamieh M, Snider DB, Sanz-Fernandez MV, Johnson JS, Keating AF, Gorden PJ, Green HB, Schoenberg KM, Baumgard LH. Intentionally induced intestinal barrier dysfunction causes inflammation, affects metabolism, and reduces productivity in lactating Holstein cows. *J Dairy Sci* 2017; 100:4113-4127.
44. Kvidera SK, Horst EA, Abuajamieh M, Mayorga EJ, Sanz-Fernandez MV, Baumgard LH. Glucose requirements of an activated immune system in lactating Holstein cows. *J Dairy Sci* 2017; 100:2360-2374.
45. Kvidera SK, Horst EA, Mayorga EJ, Sanz-Fernandez MV, Abuajamieh M, and Baumgard LH. Estimating glucose requirements of an activated immune system in growing pigs. *J Anim Sci* 2017; 95:5020-5029.
46. Kvidera SK, Horst EA, Sanz-Fernandez MV, Abuajamieh M, Ganesan S, Gorden PJ, Green HB, Schoenberg KM, Trout WE, Keating AF, Baumgard LH. Characterizing effects of feed restriction and glucagon-like peptide 2 administration on biomarkers of inflammation and intestinal morphology. *J Dairy Sci* 2017d; 100:9402-9417.
47. Lambert GP, Gisolfi CV, Berg DJ, Moseley PL, Oberley LW, Kregel KC. Selected contribution: Hyperthermia-induced intestinal permeability and the role of oxidative and nitrosative stress. *J Appl Physiol* 2002; 92:1750-1761.
48. Lanza-Jacoby S, Phetteplace H, Sedkova N, Knee G. Sequential alterations in tissue lipoprotein lipase, triglyceride secretion rates, and serum tumor necrosis factor alpha during *Escherichia coli* bacteremic sepsis in relation to the development of hypertriglyceridemia. *Shock* 1998; 9:46-51.
49. Leon LR. Heat stroke and cytokines. *Prog Brain Res* 2007; 162:481-524.
50. Lewis RS. Calcium signaling mechanisms in T lymphocytes. *Annu Rev Immunol* 2001; 19:497-521.
51. Li S, Khafipour E, Krause DO, Kroeker A, Rodriguez-Lecompte JC, Gozho GN, Plaizier JC. Effects of subacute ruminal acidosis challenges on fermentation and endotoxins in the rumen and hindgut of dairy cows. *J Dairy Sci* 2012; 95:294-303.
52. Liang H, Hussey SE, Sanchez-Avila A, Tantiwong P, Musi N. Effect of lipopolysaccharide on inflammation and insulin action in human muscle. *PLoS One* 2013; 8:e63983.
53. Lim CL, Wilson G, Brown L, Coombes JS, Mackinnon LT. Pre-existing inflammatory state compromises heat tolerance in rats exposed to heat stress. *Am J Physiol Regul Integr Comp Physiol* 2007; 292:R186-194.
54. Looor JJ, Dann HM, Everts RE, Oliveira R, Green CA, Guretzky NAJ, Rodriguez-Zas SL, Lewin HA, Drackley JK. Temporal gene expression profiling of liver from periparturient dairy cows reveals complex adaptive mechanisms in hepatic function. *Physiol Genomics* 2005; 23:217-226.
55. Malcolm DS, Zaloga GP, Holaday JW. Calcium administration increases the mortality of endotoxic shock in rats. *Crit Care Med* 1989; 17:900-903.
56. Mani V, Weber TE, Baumgard LH, Gabler NK. Growth and development symposium: endotoxin, inflammation, and intestinal function in livestock. *J Anim Sci* 2012; 90:1452-1465.
57. Maratou E, Dimitriadis G, Kollias A, Boutati E, Lambadiari V, Mitrou P, Raptis SA. Glucose transporter expression on the plasma membrane of resting and activated white blood cells. *Eur J Clin Invest* 2007; 37:282-290.
58. Martel CA, Mamedova LK, Minton JE, Jones ML, Carroll JA, Bradford BJ. Continuous low-dose infusion of tumor necrosis factor alpha in adipose tissue elevates adipose tissue interleukin 10 abundance and fails to alter metabolism in lactating dairy cows. *J Dairy Sci* 2014; 97:4897-4906.
59. McGuinness OP. Defective glucose homeostasis during infection. *Annu Rev Nutr* 2005; 25:9-35.
60. Minuti A, Ahmed S, Trevisi E, Piccioli-Cappelli F, Bertoni G, Jahan N, Bani P. Experimental acute rumen acidosis in sheep: Consequences on clinical, rumen, and gastrointestinal permeability conditions and blood chemistry. *J Anim Sci* 2014; 92:3966-3977.
61. Mullins CR, Mamedova LK, Brouk MJ, Moore CE, Green HB, Perfield KL, Smith JF, Harner JP, Bradford BJ. Effects of monensin on metabolic parameters, feeding behavior, and productivity of transition dairy cows. *J Dairy Sci* 2012; 95:1323-1336.
62. Nocek JE. Bovine acidosis: Implications on laminitis. *J Dairy Sci* 1997; 80:1005-1028.
63. O'Boyle NJ, Contreras GA, Mattmiller SA, Sordillo LM. Changes in glucose transporter expression in monocytes of periparturient dairy cows. *J Dairy Sci* 2012; 95:5709-5719.
64. Piantoni P, Abeyta MA, Schroeder GF, Ramirez-Ramirez HA, Tucker HA, Baumgard LH. Induction of leaky gut through feed restriction or abomasal infusion of resistant starch in healthy post-peak lactating cows. *J Dairy Sci* 2018; (Suppl. 2): 228.
65. Palsson-McDermott EM, O'Neill LA. The Warburg effect then and now: from cancer to inflammatory diseases. *Bioessays* 2013; 35:965-973.
66. Pearce SC, Mani V, Weber TE, Rhoads RP, Patience JF, Baumgard LH, Gabler NK. Heat stress and reduced plane of nutrition decreases intestinal integrity and function in pigs. *J Anim Sci* 2013; 91:5183-5193.
67. Poggi M, Bastelica D, Gual P, Iglesias MA, Gremaux T, Knauf C, Peiretti F, Verdier M, Juhan-Vague I, Tanti JF, Burcelin R, Alessi MC. C3H/HeJ mice carrying a toll-like receptor 4 mutation are protected against the development of insulin resistance in white adipose tissue in response to a high-fat diet. *Diabetologia* 2007; 50:1267-1276.
68. Rhoads ML, Rhoads RP, VanBaale MJ, Collier RJ, Sanders SR, Weber WJ, Crooker BA, Baumgard LH. Effects of heat stress and plane of nutrition on lactating Holstein cows: I. Production, metabolism, and aspects of circulating somatotropin. *J Dairy Sci* 2009; 92:1986-1997.
69. Rodiño-Janeiro B K, Alonso-Cotoner C, Pigrau M, Lobo B, Vicario M, Santos J. Role of corticotropin-releasing factor in gastrointestinal permeability. *J Neurogastroenterol Motil* 2015; 21:33-50.
70. Rollwagen FM, Madhavan S, Singh A, Li YY, Wolcott K, Maheshwari R. IL-6 protects enterocytes from hypoxia-induced apoptosis by induction of bcl-2 mRNA and reduction of fas mRNA. *Biochem Biophys Res Commun* 2006; 347:1094-1098.
71. Santos J, Benjamin J, Yang PC, Prior T, Perdue MH. Chronic stress impairs rat growth and jejunal epithelial barrier function: role of mast cells. *Am J Physiol Gastrointest Liver Physiol* 2000; 278: G847-G854.
72. Sanz-Fernandez MV, Pearce SC, Gabler NK, Patience JF, Wilson ME, Socha MT, Torrison JL, Rhoads RP, Baumgard LH. Effects of supplemental zinc amino acid complex on gut integrity in heat-stressed growing pigs. *Animal* 2014; 8:43-50.
73. Skarnes RC, Chedid LC. Biological degradation and inactivation of endotoxin (chromate-labeled). In: Landy M, Braun W, eds. *Bacterial Endotoxins*. New Brunswick, NJ: Rutgers University Press, 1964; 575-587.
74. Smith F, Clark JE, Overman BL, Tozel CC, Huang JH, Rivier JEF, Blikslager AT, Moeser AJ. Early weaning stress impairs development of mucosal barrier function in the porcine intestine. *Am J Physiol Gastrointest Liver Physiol* 2010; 298: G352-G363.
75. Spitzer JA, Nelson KM, Fish RE. Time course of changes in gluconeogenesis from various precursors in chronically endotoxemic rats. *Metabolism* 1985; 34:842-849.
76. Tennant B, Reina-Guerra M, Harrold D. Metabolic response of calves following acute experimental endotoxemia. *Ann Rech Veter* 1973; 4:135-147.
77. Toribio RE, Kohn CW, Hardy J, Rosol TJ. Alterations in serum parathyroid hormone and electrolyte concentrations and urinary excretion of electrolytes in horses with induced endotoxemia. *J Vet Intern Med* 2005; 19:223-231.
78. Tough DF, Sun S, Sprent J. T cell stimulation in vivo by lipopolysaccharide (LPS). *J Exp Med* 1997; 185:2089-2094.

79. Vanuytsel T, van Wanrooy S, Vanheel H, Vanormelingen C, Verschuere S, Houben E, Salim Rasool S, Tóth J, Holvoet L, Farré R, Van Oudenhove L, Boeckxstaens G, Verbeke K, Tack J. Psychological stress and corticotropin-releasing hormone increase intestinal permeability in humans by a mast cell-dependent mechanism. *Gut* 2014; 63:1293-1299.
80. Waldron MR, Nonnecke BJ, Nishida T, Horst RL, Overton TR. Effect of lipopolysaccharide infusion on serum macromineral and vitamin D concentrations in dairy cows. *J Dairy Sci* 2003; 86:3440-3446
81. Waldron MR, Nishida T, Nonnecke BJ, Overton TR. Effect of lipopolysaccharide on indices of peripheral and hepatic metabolism in lactating cows. *J Dairy Sci* 2003b; 86:3447-3459.
82. Waldron MR, Kulick AE, Bell AW, Overton TR. Acute experimental mastitis is not causal toward the development of energy-related metabolic disorders in early postpartum dairy cows. *J Dairy Sci* 2006; 89:596-610.
83. Wallon C, Yang PC, Keita AV, Ericson AC, McKay DM, Sherman PM, Perdue MH, Söderholm JD. Corticotropin-releasing hormone (CRH) regulates macromolecular permeability via mast cells in normal human colonic biopsies in vitro. *Gut* 2008; 57:50-58.
84. Wannemacher RW, Beall FA, Canonico PG, Dinterman RE, Hadick CL, Neufeld HA. Glucose and alanine metabolism during bacterial infections in rats and rhesus monkeys. *Metabolism* 1980; 29:201-212.
85. Yuan K, Farney JK, Mamedova LK, Sordillo LM, Bradford BJ. TNF α Altered Inflammatory Responses, Impaired Health and Productivity, but Did Not Affect Glucose or Lipid Metabolism in Early-Lactation Dairy Cows. *PLoS One* 2013; e80316.
86. Zarrin M, Wellnitz O, van Dorland HA, Gross JJ, Bruckmaier RM. Hyperketonemia during lipopolysaccharide-induced mastitis affects systemic and local intramammary metabolism in dairy cows. *J Dairy Sci* 2014; 97:3531-3541.