

Uterine health problems – Risk factors and effects on herd performance

Mark J. Thomas, DVM, DABVP; Matias L. Stangaferro, DVM, MS, PhD; Rita Couto Serrenho, DVM, MSc
Dairy Health and Management Services, Lowville, NY 13367

Abstract

A successful calving event and the following transition period are critically important to the future reproductive performance and productivity of the dairy cow. Although the diagnosis and treatment of uterine diseases are generally considered straightforward by most veterinarians, there are many misconceptions and a lack of data to support many of the recommended treatments. Other metabolic disorders such as hypocalcemia and hyperketonemia can further complicate the diagnosis and treatment plan.

Not only is an understanding of reproductive physiology and normal uterine involution important, but also the economic outcomes of uterine disease. The effects on milk production, fertility, risk to other diseases, and exit from the herd need to be recognized by both the dairy herd managers and veterinarians alike.

This article will review the definitions, risk factors, treatment options and economics of uterine diseases: retained placenta (RP), metritis, and endometritis.

Key words: uterine health, metritis, transition cow disease

Résumé

Un vêlage réussi et la période de transition suivante sont très importants pour les performances reproductives et la productivité futures de la vache laitière. Bien que le diagnostic et le traitement des maladies utérines soient généralement considérés simples par la plupart des vétérinaires, il existe plusieurs idées fausses et un manque de données pour supporter plusieurs des traitements recommandés. D'autres troubles métaboliques comme l'hypocalcémie et l'acétonémie tendent à compliquer davantage le diagnostic et le plan de traitement.

La connaissance de la physiologie de la reproduction et de l'involution utérine normale importe autant que la reconnaissance des résultats économiques de la maladie utérine. Les effets sur la production laitière, la fertilité, le risque de développer d'autres maladies et le retrait du troupeau doivent être reconnus autant par les gestionnaires de la ferme laitière que par les vétérinaires.

Cet article fait le point sur les définitions, les facteurs de risque, les options de traitement et les retombées économiques des maladies utérines : rétention placentaire, métrite et endométrite.

Introduction

High milk yield and reduced involuntary culling have been shown to improve overall herd profitability.^{4,18} Uterine disease decreases milk yield,^{a,20} fertility,⁵ and herd survival.^{12,14} Thus, it is expected that RP and metritis can have a major impact on the economics of a dairy operation. Understanding the risk factors and the prevention of uterine disease, along with proper diagnosis and treatment choices, is therefore important.

The transition period is generally defined as 3 weeks prior to 3 weeks following calving. This period has the greatest risk of metabolic disease, with some herds having 30 to 50% of transition cows affected by 1 or more diseases.¹⁹ There are many risk factors for RP, metritis, and endometritis,²⁴ which, in turn, are risk factors for other metabolic diseases such as hypercalcemia, hyperketonemia, and displaced abomasum (DA).⁷ Uterine damage from dystocia and more importantly, nutritional issues during the pre-calving close-up period, are major areas of focus for the prevention of disease.

Risk Factors

One important aspect in the discussion of uterine disease is the difference between contamination and infection of the uterus. Uterine contamination does not always lead to metritis or endometritis.³⁰ Uterine infections are generally considered non-specific in nature and are caused by environmental bacteria.¹

Delayed uterine involution is the factor most commonly associated with uterine infection. Any disruption of normal calving such as dystocia, twins and stillbirth, increase the risk of delayed involution¹⁵ and therefore can lead to infection.

Metabolic diseases such as hypocalcemia, hyperketonemia, and DA are also associated with an increased risk of uterine disease. The exact mechanism underlying this association has not been determined, but a failure of the immune system is believed to play a part in this increased risk.³

Diagnosis

The diagnostic definition of RP is straightforward; visible fetal membranes greater than 24 hours after calving. The diagnosis of clinical metritis can be more challenging and is a current topic of discussion, but is defined as a fetid,

watery uterine discharge often accompanied by fever and/or systemic signs of disease.³⁰ Clinical endometritis is defined as the presence of pus in the vagina > 21 days post-calving. Clinical endometritis must be differentiated from cases of vaginitis that can also present with a purulent discharge. This distinction can be difficult given that common methods of diagnosis (e.g. Metricek device or gloved hand) cannot differentiate between vaginitis and endometritis.²⁶

Treatment

Treatment protocols need to be based on an accurate diagnosis of the uterine disease along with any concurrent metabolic disorders. A quote from Frazer provides the current thinking on various treatment modalities: "This postpartum metritis-delayed uterine involution syndrome is extremely frustrating for a veterinary clinician to manage since there is no scientifically proven protocol that will enhance uterine contraction and promote evacuation of the fetid uterine contents. Supportive measures (anti-inflammatory medication and systemic antibiotics) may help to maintain the cow's appetite and rumen motility, but this author remains unconvinced that any current hormonal therapy actually works."⁸

A detailed review of the treatment of RP, metritis, and endometritis can be found in the literature.⁵ The current data does not support the use of prostaglandins for the treatment of RP or metritis.^{6,11,21} Additionally, infusions of intrauterine disinfectants and/or antibiotics have not shown a benefit in terms of future fertility and may in fact be detrimental.⁹ These treatments are also extra-label in the USA and present the risk for antibiotic residues.¹³ Non-traditional treatments such as intra-uterine infusion of mannose or bacteriophage²³ or dextrose^{2,23} have not been shown to be effective for the treatment of uterine infections.

For severe, acute toxic metritis, systemic antibiotics, non-steroidal anti-inflammatory drugs (NSAIDS), supportive care, and treatment of concurrent metabolic disease represent the accepted treatment regimen. The antibiotic choice should be based on spectrum of activity, withholding times and economics with the knowledge that *E. coli*, *T. pyogenes*, and *F. necrophorum* are the most common isolates from metritic cows.³⁰ NSAIDS are generally used based on first-principle concepts for treatment of pyrexia, as literature support for such use appears to be lacking^{5,16,22,27} and the routine use of flunixin meglumine in peri-parturient cows has been shown to increase the prevalence of RP and metritis.²⁵

Prevention

Given the multifactorial pathophysiology of uterine diseases, prevention of uterine contamination and infection needs to be focused on overall transition cow management to minimize predisposing factors.³¹ The role of nutrition in preventing metabolic disorders such as hyperketonemia,

hypocalcemia, and DA is of major importance. Attention to maternity area hygiene is another important area for the reduction of uterine contamination at calving.²⁸ Training of maternity pen caregivers to ensure proper assistance at birthing, dystocia management and hygiene are other important considerations.^{17,28,29}

The use of vaccines targeting uterine infection pathogens has been investigated²³ and holds potential promise for future commercial use. Immune modulators such as pegbovigrastim have not been shown to be beneficial for the prevention of uterine disease.³²

Economics

There are a number of economic models that have been used to calculate the cost of transition cow diseases.^{3,10,14,20} It is interesting to note that these models all calculate a very similar economic loss from uterine disease with an average cost per case for RP/metritis combined of \$322 and \$576 per case for primiparous and multiparous cows, respectively. Milk loss represents the largest component of the cost of metritis, with choice of treatment having a minimal effect on overall cost per case.²⁰

Although these costs represent real economic losses, the overall low incidence of these diseases in most well-managed herds does not represent a significant economic return in the reduction of incidence. The goal is to always prevent disease, but the savings from reducing RP/metritis by 1% in a 1000-cow herd is less than \$3000/year.^a Therefore, prioritization of disease prevention strategies must be considered individually for each dairy and each disease.

Conclusions

High milk production and cow longevity are important factors for overall dairy farm profitability. Management of the nutritional and environmental needs of the dairy cow along with proper training of employees in maternity care are critically important to reduce the risk of postpartum disease and ensure a successful transition period. Transition cow diseases are multifactorial and interrelated. Attention to the prevention, correct diagnosis and effective treatment of uterine diseases can result in positive economic returns, but this must be prioritized with other management and prevention strategies that can perhaps yield greater returns.

Endnote

^a Thomas MJ, Stangaferro ML. Unpublished data, 2019.

References

1. Bicalho RC, Machado VS, Bicalho MLS, Gilbert RO, Teixeira AGV, Caixeta LS, Pereira RVV. Molecular and epidemiological characterization of bovine intrauterine *Escherichia coli*. *J Dairy Sci* 2010;93:5818-5830.

2. Brick TA, Schuenemann GM, Bas S, Daniels JB, Pinto CR, Rings DM, Rajala-Schultz PJ. Effect of intrauterine dextrose or antibiotic therapy on reproductive performance of lactating dairy cows diagnosed with clinical endometritis. *J Dairy Sci* 2012;95:1894-1905.
3. Correa MT, Erb H, Scarlett J. Path analysis for seven postpartum disorders of Holstein cows. *J Dairy Sci* 1993;76:1305-1312.
4. De Vries A. Reproduction: Economics of reproductive efficiency. *Revista Colombiana de Ciencias Pecuarias* 2017;30:218-221.
5. Drillich M, Beetz O, Pfutzner A, Sabin M, Sabin HJ, Kutzer P, Nattermann H, Heuwieser W. Evaluation of a systemic antibiotic treatment of toxic puerperal metritis in dairy cows. *J Dairy Sci* 2001;84:2010-2017.
6. Dubuc J, Duffield TF, Leslie KE, Walton JS, LeBlanc SJ. Randomized clinical trial of antibiotic and prostaglandin treatments for uterine health and reproductive performance in dairy cows. *J Dairy Sci* 2011;94:1325-1338.
7. Erb HN, Martin SW, Ison N, Swaminathan S. Interrelationships between production and reproductive diseases in Holstein cows. Path analysis. *J Dairy Sci* 1981;64:282-289.
8. Frazer GS. A rational basis for therapy in the sick postpartum cow. *Vet Clin North Am Food Anim Pract* 2005;21:523-568.
9. Fuquay JW, Harris RA, McGee WH, Beatty JF, Arnold BL. Routine postpartum treatment of dairy cattle with intrauterine neomycin sulfate boluses. *J Dairy Sci* 1975;58:1367-1369.
10. Galligan D. Economic assessment of animal health performance. *Vet Clin North Am Food Anim Pract* 2006;22:207-227.
11. Galvao KN, Frajblat M, Brittin SB, Butler WR, Guard CL, Gilbert RO. Effect of prostaglandin F2 α on subclinical endometritis and fertility in dairy cows. *J Dairy Sci* 2009;92:4906-4913.
12. Gilbert RO, Shin ST, Guard CL, Erb HN, Frajblat M. Incidence of endometritis and its effects on reproductive performance of dairy cows. *Theriogenology* 2005;64:1879-1888.
13. Gorden PJ, Ydstie JA, Kleinhenz MD, Wulf LW, Gehring R, Lee CJ, Wang C, Coetzee JF. A study to examine the relationship between metritis severity and depletion of oxytetracycline in plasma and milk after intrauterine infusion. *J Dairy Sci* 2016;99:8314-8322.
14. Guard D. The costs of common diseases of dairy cattle, in *Proceedings. CVC* in San Diego, 2008.
15. Hussain AM, Daniel RCW, O'Boyle D. Postpartum uterine flora following normal and abnormal puerperium in cows. *Theriogenology* 1990;34:291-302.
16. Jeremejeva J, Orro T, Waldmann A. Treatment of dairy cows with PGF2 α or NSAID, in combination with antibiotics, in cases of postpartum uterine inflammation. *Acta Vet Scand* 2012;54:45.
17. Kristula MA, Smith BI. Care report—Veterinary specific training to manage dairy cows at calving time. *Bov Pract* 2011;45:78-86.
18. Krpálková L, Cabrera VE, Kvapilík J, Burdych J, Crump P. Associations between age at first calving, rearing average daily weight gain, herd milk yield and dairy herd production, reproduction, and profitability. *J Dairy Sci* 2014;97:6573-6582.
19. LeBlanc SJ, Duffield TF, Leslie KE, Bateman KG, Keefe GP, Walton JS, Johnson WH. Defining and diagnosing postpartum clinical endometritis and its impact on reproductive performance in dairy cows. *J Dairy Sci* 2002;85:2223-2236.
20. Liang D, Arnold LM, Stowe CJ, Harmon RJ, Bewley JM. Estimating US dairy clinical disease costs with a stochastic simulation model. *J Dairy Sci* 2017;100:1472-1486.
21. Lima FS, Vieira-Neto A, Vasconcellos GSFM, Mingoti RD, Karakaya E, Solé E, Bisinotto RS, Martinez N, Risco CA, Galvao KN, Santos JEP. Efficacy of ampicillin trihydrate or ceftiofur hydrochloride for treatment of metritis and subsequent fertility in dairy cows. *J Dairy Sci* 2014;97:5401-5414.
22. Lomb J, Neave HW, Weary DM, LeBlanc SJ, Huzzey JM, Von Keyserlingk MAG. Changes in feeding, social, and lying behaviors in dairy cows with metritis following treatment with a nonsteroidal anti-inflammatory drug as adjunctive treatment to an antimicrobial. *J Dairy Sci* 2018;101:4400-4411.
23. Machado VS, Bicalho MLdS, Meira Jr EBdS, Rossi R, Ribeiro BL, Lima S, Santos T, Kussler A, Foditsch C, Ganda EK, Oikonomou G, Cheong SH, Gilbert RO, Bicalho RC. Subcutaneous immunization with inactivated bacterial components and purified protein of *Escherichia coli*, *Fusobacterium necrophorum* and *Trueperella pyogenes* prevents puerperal metritis in Holstein dairy cows. *PLoS ONE* 2014;9:e1734.
24. McNaughton AP, Murray RD. Structure and function of the bovine fetomaternal unit in relation to the causes of retained fetal membranes. *Vet Rec* 2009;165:615-622.
25. Newby NC, Leslie KE, Dingwell HP, Kelton DF, Weary DM, Neuder L, Duffield TF. The effects of periparturient administration of flunixin meglumine on the health and production of dairy cattle. *J Dairy Sci* 2017;100:582-587.
26. Pleticha S, Drillich M, Heuwieser W. Evaluation of the Metricheck device and the gloved hand for the diagnosis of clinical endometritis in dairy cows. *J Dairy Sci* 2009;92:5429-5435.
27. Pohl A, Bertulat S, Borchardt S, Burfeind O, Heuwieser W. Randomized, controlled clinical trial on the efficacy of nonsteroidal antiinflammatory drugs for the treatment of acute puerperal metritis in dairy cows. *J Dairy Sci* 2016;99:8241-8249.
28. Schuenemann GM, Nieto I, Bas S, Galvao KN, Workman J. Dairy calving management: Effect of perineal hygiene scores on metritis. *J Dairy Sci* 2011;94(E-Suppl. 1):744.
29. Schuenemann GM, Bas S, Workman JD, Rajala-Schultz R. Dairy reproductive management: Assessing a comprehensive continuing education program for veterinary practitioners. *J Vet Med Educ* 2010;37:289-298.
30. Sheldon IM, Williams EJ, Miller AN, Nash DM, Herath S. Uterine diseases in cattle after parturition. *Vet J* 2008;176:115-121.
31. Stevenson JS, Call EP. Reproductive disorders in the periparturient dairy cow. *J Dairy Sci* 1988;71:2572-2583.
32. Zinicola M, Korzec H, Teixeira AGV, Ganda EK, Bringhenti L, Tomazi ACCH, Gilbert RO, Bicalho RC. Effects of pegbovigrastim administration on periparturient diseases, milk production, and reproductive performance of Holstein cows. *J Dairy Sci* 2018;101:11199-11217.