





ZACTRAN[®]

(gamithromycin)

150 mg/mL ANTIMICROBIAL

NADA 141-328, Approved by FDA

For subcutaneous injection in beef and non-lactating dairy cattle only.
Not for use in female dairy cattle 20 months of age or older or in calves to be processed for veal.

Caution: Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

READ ENTIRE BROCHURE CAREFULLY BEFORE USING THIS PRODUCT.

INDICATIONS

ZACTRAN is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis* in beef and non-lactating dairy cattle. ZACTRAN is also indicated for the control of respiratory disease in beef and non-lactating dairy cattle at high risk of developing BRD associated with *Mannheimia haemolytica* and *Pasteurella multocida*.

CONTRAINDICATIONS

As with all drugs, the use of ZACTRAN is contraindicated in animals previously found to be hypersensitive to this drug.

WARNING: FOR USE IN CATTLE ONLY. NOT FOR USE IN HUMANS. KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN. NOT FOR USE IN CHICKENS OR TURKEYS.

The material safety data sheet (MSDS) contains more detailed occupational safety information. To report adverse effects, obtain an MSDS or for assistance, contact Merial at 1-888-637-4251.

RESIDUE WARNINGS: Do not treat cattle within 35 days of slaughter. Because a discard time in milk has not been established, do not use in female dairy cattle 20 months of age or older. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

PRECAUTIONS

The effects of ZACTRAN on bovine reproductive performance, pregnancy, and lactation have not been determined. Subcutaneous injection of ZACTRAN may cause a transient local tissue reaction in some cattle that may result in trim loss of edible tissues at slaughter.

ADVERSE REACTIONS

Transient animal discomfort and mild to moderate injection site swelling may be seen in cattle treated with ZACTRAN.

EFFECTIVENESS

The effectiveness of ZACTRAN for the treatment of BRD associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* was demonstrated in a field study conducted at four geographic locations in the United States. A total of 497 cattle exhibiting clinical signs of BRD were enrolled in the study. Cattle were administered ZACTRAN (6 mg/kg BW) or an equivalent volume of sterile saline as a subcutaneous injection once on Day 0. Cattle were observed daily for clinical signs of BRD and were evaluated for clinical success on Day 10. The percentage of successes in cattle treated with ZACTRAN (58%) was statistically significantly higher ($p < 0.05$) than the percentage of successes in the cattle treated with saline (19%).

The effectiveness of ZACTRAN for the treatment of BRD associated with *M. bovis* was demonstrated independently at two U.S. study sites. A total of 502 cattle exhibiting clinical signs of BRD were enrolled in the studies. Cattle were administered ZACTRAN (6 mg/kg BW) or an equivalent volume of sterile saline as a subcutaneous injection once on Day 0. At each site, the percentage of successes in cattle treated with ZACTRAN on Day 10 was statistically significantly higher than the percentage of successes in the cattle treated with saline (74.4% vs. 24% [$p < 0.001$], and 67.4% vs. 46.2% [$p = 0.002$]). In addition, in the group of calves treated with gamithromycin that were confirmed positive for *M. bovis* (pre-treatment nasopharyngeal swabs), there were more calves at each site (45 of 57 calves, and 5 of 6 calves) classified as successes than as failures.

The effectiveness of ZACTRAN for the control of respiratory disease in cattle at high risk of developing BRD associated with *Mannheimia haemolytica* and *Pasteurella multocida* was demonstrated in two independent studies conducted in the United States. A total of 467 crossbred beef cattle at high risk of developing BRD were enrolled in the study. ZACTRAN (6 mg/kg BW) or an equivalent volume of sterile saline was administered as a single subcutaneous injection within one day after arrival. Cattle were observed daily for clinical signs of BRD and were evaluated for clinical success on Day 10 post-treatment. In each of the two studies, the percentage of successes in the cattle treated with ZACTRAN (86% and 78%) was statistically significantly higher ($p = 0.0019$ and $p = 0.0016$) than the percentage of successes in the cattle treated with saline (36% and 58%).

Marketed by Merial Limited

3239 Satellite Blvd., Duluth, GA 30096-4640 U.S.A.
Made in Austria

©ZACTRAN is a registered trademark of Merial Limited.
©2012 Merial Limited. All rights reserved. Rev. 03/2012

TAKE THE STRESS OUT OF BRD.

Target high-risk calves with ZACTRAN® (gamithromycin).

EXZACTLY THE RIGHT ANSWER FOR BRD.

With rising feed costs and tight margins, your clients are as stressed as their long-haul cattle. That's why they need ZACTRAN.

ZACTRAN delivers rapid onset¹ and 10-day duration² against the most prevalent causes of BRD in a single dose.^{3,4} And most cattle stayed healthy with ZACTRAN, meaning fewer

retreatments.⁵ Talk to your clients about prescription ZACTRAN. It's exZACTLY what you need to help them control BRD risk with one treatment.



Give subcutaneously at 2 mL/110 lbs.

ZACTRAN[®]
(gamithromycin)

IMPORTANT SAFETY INFORMATION: For use in cattle only. Do not treat cattle within 35 days of slaughter. Because a discard time in milk has not been established, do not use in female dairy cattle 20 months of age or older, or in calves to be processed for veal. The effects of ZACTRAN on bovine reproductive performance, pregnancy and lactation have not been determined.

¹ Siffman RL, Wolff WA, Holste JE, et al. Field efficacy evaluation of gamithromycin for treatment of bovine respiratory disease in cattle at feedlots. *Intern J Appl Res Vet Med.* 2011;9(2):171-180.

² Lechtenberg K, Daniels CS, Royer GC, et al. Field efficacy study of gamithromycin for the control of bovine respiratory disease in cattle at high risk of developing the disease. *Intern J Appl Res Vet Med.* 2011;9(2):189-197.

³ ZACTRAN product label.

⁴ Kahn, CM. *Merck Veterinary Manual*. 10th edition. 2010:1319.

⁵ Van Donkersgoed J, Merrill JK. A comparison of tilmosin to gamithromycin for on-arrival treatment of bovine respiratory disease in feeder steers. *Bovine Practitioner.* 2012;46(1):46-51.



©ZACTRAN is a registered trademark of Merial.
©2014 Merial, Inc., Duluth, GA. All rights reserved.
RUMIOTD1301-B (04/14)