



Vol. 48, No. 2  
SUMMER 2014

# *The Bovine* **PRACTITIONER**

THE OFFICIAL PUBLICATION OF THE  
AMERICAN ASSOCIATION OF BOVINE PRACTITIONERS



© Copyright American Association of Bovine Practitioners; open access distribution.

## **47th ANNUAL CONFERENCE**

American Association of Bovine Practitioners  
Albuquerque, New Mexico  
September 18-20, 2014

Property of  
**Medical Sciences Library**  
Texas A&M University


# Moraxella bovoculi Problems?

# Moraxella bovoculi SOLUTION!


## The Addison Autogenous Program

- **Complete Diagnostic and Autogenous Support**
- **Fast Delivery on Initial Order**
- **Excellent Syringeability**
- **Exceptional Safety Record**


MAXI/GUARD® Pinkeye Bacterin has earned the trust of bovine practitioners as the preferred pinkeye preventative when the disease is caused solely by *M. bovis*. However, the problem can be caused by the combination of *M. bovis* and *M. bovoculi* bacteria. Unfortunately no commercial vaccines provide that coverage. When *M. bovoculi* is present, your best option is utilizing MAXI/GUARD® along with the Addison Autogenous Program. We can assist you from diagnosing the problem to creating the solution!



© Copyright American Association of Bovine Practitioners; open access distribution.



*From the LEADERS in pinkeye prevention!*



Addison Biological Laboratory, Inc.

[www.addisonlabs.com](http://www.addisonlabs.com)

800-331-2530

 /AddisonLabs





# The Bovine PRACTITIONER

**EDITOR:**  
**ROBERT A. SMITH, DVM**  
3404 Live Oak Lane  
Stillwater, Oklahoma 74075  
Tel: (405) 372-8666  
Fax: (405) 743-8422

**PUBLISHER:**  
**VM PUBLISHING COMPANY**  
205 W. 7th Avenue, Suite 201G  
Stillwater, Oklahoma 74074  
Tel/Fax: (405) 533-1883  
kellijo.vmpubco@gmail.com

## OFFICERS

|                                       |  |
|---------------------------------------|--|
| <b>President</b> .....                | <b>Daniel L. Grooms, DVM</b><br>Williamston, Michigan  |
| <b>President-Elect</b> .....          | <b>John M. Davidson, DVM</b><br>College Station, Texas |
| <b>Vice President</b> .....           | <b>K. Fred Gingrich II, DVM</b><br>Ashland, Ohio       |
| <b>Past President</b> .....           | <b>Nigel Cook, MRCVS</b><br>Madison, Wisconsin         |
| <b>Executive Vice President</b> ..... | <b>M. Gatz Riddell, Jr., DVM</b><br>Auburn, Alabama    |
| <b>Treasurer</b> .....                | <b>Brian Reed, DVM</b><br>Lititz, Pennsylvania         |
| <b>Parliamentarian</b> .....          | <b>Patty Scharko, DVM</b><br>Columbia, South Carolina  |

## BOARD OF DIRECTORS

**Mark J. Thomas, DVM**  
Carthage, New York  
1st District, 2014

**Thomas B. Massie, Jr., DVM**  
Washington, Virginia  
2nd District, 2015

**James A. Brett, DVM**  
Starkville, Mississippi  
3rd District, 2016

**Garold H. Koester, DVM**  
Cadillac, Michigan  
4th District, 2014

**Hunter Lang, DVM**  
Prairie du Sac, Wisconsin  
5th District, 2015

**Sarah Overby, DVM**  
Freeport, Minnesota  
6th District, 2016

**Rodney K. Chapman, DVM**  
Versailles, Missouri  
7th District, 2014

**Joe M. Hillhouse, DVM**  
Panhandle, Texas  
8th District, 2015

**Dee Griffin, DVM**  
Clay Center, Nebraska  
9th District, 2016

**Stephen A. Smalley, VMD**  
Chandler, Arizona  
10th District, 2014

**Frederick J. Muller, DVM**  
Sunnyside, Washington  
11th District, 2015

**Wayne S. Shewfelt, DVM**  
Tavistock, Ontario, Canada  
12th District, 2016

**Eric Behlke, DVM**  
Okotoks, Alberta, Canada  
13th District, 2014

### AVMA DELEGATE:

**Brian J. Gerloff, DVM**  
Marengo, Illinois

### EXHIBITS CHAIRMAN:

**Charles W. Hatcher, DVM**  
College Grove, Tennessee

### ADVERTISING AGENCY:

**J. T. O'Brien & Assoc.**  
12118 Nieman Road  
Overland Park, Kansas 66213  
(913) 579-4084 (cell)

THE BOVINE PRACTITIONER is published by The American Association of Bovine Practitioners, Inc., at **VM Publishing Co., 205 W. 7th, Suite 201G, Stillwater, OK 74074 (Telephone and Fax: 405-533-1883)**. Over 6000 copies are mailed to veterinarians engaged in cattle practice and allied institutions. Send new address (please include old address) to:

**AABP, P.O. Box 3610, Auburn, AL 36831-3610**

**Tel: 1-800-269-2227; FAX: 334-821-9532**

**e-mail: aabphq@aabp.org**

**http://www.aabp.org**

THE BOVINE PRACTITIONER is the official publication of The American Association of Bovine Practitioners, published in the spring and summer annually. It also serves as a communication medium between bovine practitioner organizations around the world. All manuscripts and communications must be presented in English.

Copies of THE BOVINE PRACTITIONER are available to non-members for \$30.00 per copy for domestic destinations, or \$35.00 for international addresses. **Contact editor for details.**

# Table of Contents

|  |     |
|--|-----|
| Estimated prevalence of lameness in 53 Northwest US dairy herds<br><i>Alea C. Hoffman, John R. Wenz, Jorge Vanegas, Dale A. Moore</i> .....  | 89  |
| Diverse outcomes of bovine viral diarrhea virus (BVDV) infections in a herd naturally infected during pregnancy—<br>a case study<br><i>Robert W. Fulton, Grant B. Rezabek, Ryan Grant, Julia F. Ridpath, Lurinda J. Burge</i> .....  | 95  |
| Comparative efficacy of tilmicosin, florfenicol, and florfenicol-flunixin meglumine for treatment of<br>undifferentiated fever in backgrounded winter-placed feedlot calves given tilmicosin metaphylactically<br>on arrival<br><i>Joyce Van Donkersgoed, John K. Merrill, Steve Hendrick</i> .....  | 103 |
| Effects of injectable fat-soluble vitamins and selenium on serum status in young beef calves<br><i>Carl R. Dahlen, Rob L. Stuart, Bryan W. Neville</i> .....   | 109 |
| Effectiveness of tildipirosin and tulathromycin in the control of bovine respiratory disease in high-risk<br>beef heifers<br><i>Riccardo Compiani, Gianluca Baldi, Maurizio Bonfanti, Daniele Fucci, Giuliano Pisoni, Stefano Jottini,<br/>Siddhartha Torres</i> .....   | 114 |
| Effects of timing of chlortetracycline in combination with decoquinatate on growth performance, health, and<br>carcass characteristics of feeder steers<br><i>Daniel U. Thomson, R. Spencer Swingle, Mark Branine, Steven J. Bartle, David A. Yates</i> .....  | 120 |
| Case study: Challenges in diagnosing hydrocyanic acid toxicity in cattle<br><i>Thomas Hairgrove, Ron Gill, Larry Redmon, Monte Rouquette, Gary Warner</i> .....  | 129 |
| Ergot toxicosis causing death in weaned beef calves<br><i>Bruce Leuschen, Steve Ensley, Paul Plummer</i> .....   | 134 |
| Comparison of milk production changes in adult Holstein cows vaccinated with a 5-way modified-live virus vaccine<br>containing a <i>Mannheimia haemolytica</i> toxoid versus a 5-way modified-live virus vaccine containing<br>a 5-way <i>Leptospira</i> bacterin<br><i>Michael D. Kleinhenz, Brian E. Miller, Sjoert Zuidhof, Patrick J. Gorden</i> ..... | 139 |
| Survival of dairy cattle presumptively suffering from idiopathic pericardial hemorrhage using parenteral<br>dexamethasone or isofluprednone acetate<br><i>Jerome Gaska, Sheila M. McGuirk, Simon F. Peek</i> .....   | 143 |
| The Michigan State University Food Systems Fellowship Program: Meeting the needs of animal agriculture<br>now and in the future<br><i>Chelsea Render, Heather Roney, Daniel Grooms</i> .....   | 148 |

## Advertisers

|  |                               |
|--|-------------------------------|
| Addison Biological Laboratory, Inc. .... | inside front cover            |
| Elanco Animal Health.....                | 99-102                        |
| Merck Animal Health .....                | front of book                 |
| Norbrook Laboratories Limited.....       | inside back cover, back cover |

*Cover photo courtesy of Dr. Pam Ruegg, Madison, Wisconsin.*

*The abstracts were translated to French by Guy Beauchamp  
and revised by Dr. Emile Bouchard.*

### Notice to Readers

All statements, opinions and conclusions contained in articles in *The Bovine Practitioner* are those of the author(s), and are not necessarily those of the American Association of Bovine Practitioners (AABP) unless specifically approved by the AABP Board of Directors.

**PRODUCT INFORMATION**

NADA 141-334, Approved by FDA.

**ZUPREVO™ 18%**  
(tildipirosin)

**Injectable Solution for Cattle****ANTIMICROBIAL DRUG**

180 mg of tildipirosin/mL. For subcutaneous injection in beef and non-lactating dairy cattle only.

**Not for use in female dairy cattle 20 months of age or older or in calves to be processed for veal.**

**CAUTION:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

**BRIEF SUMMARY:** for full prescribing information use package insert.

**INDICATIONS:** Zuprevo™ 18% is indicated for the treatment of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni* in beef and non-lactating dairy cattle, and for the control of respiratory disease in beef and non-lactating dairy cattle at high risk of developing BRD associated with *M. haemolytica*, *P. multocida*, and *H. somni*.

**WARNINGS: FOR USE IN ANIMALS ONLY. NOT FOR HUMAN USE. KEEP OUT OF REACH OF CHILDREN. TO AVOID ACCIDENTAL INJECTION, DO NOT USE IN AUTOMATICALLY POWERED SYRINGES WHICH HAVE NO ADDITIONAL PROTECTION SYSTEM. IN CASE OF HUMAN INJECTION, SEEK MEDICAL ADVICE IMMEDIATELY AND SHOW THE PACKAGE INSERT OR LABEL TO THE PHYSICIAN.**

Avoid direct contact with skin and eyes. If accidental eye exposure occurs, rinse eyes with clean water. If accidental skin exposure occurs, wash the skin immediately with soap and water. Tildipirosin may cause sensitization by skin contact.

For technical assistance or to report a suspected adverse reaction, call: 1-800-219-9286.

For customer service or to request a Material Safety Data Sheet (MSDS), call: 1-800-211-3573. For additional Zuprevo 18% information go to [www.zuprevo.com](http://www.zuprevo.com).

For a complete listing of adverse reactions for Zuprevo 18% reported to CVM see: <http://www.fda.gov/AnimalVeterinary/SafetyHealth>.

**DO NOT USE ZUPREVO 18% IN SWINE.**

Fatal adverse events have been reported following the use of tildipirosin in swine. NOT FOR USE IN CHICKENS OR TURKEYS.

**RESIDUE WARNING:** Cattle intended for human consumption must not be slaughtered within 21 days of the last treatment. Do not use in female dairy cattle 20 months of age or older. Use of this drug product in these cattle may cause milk residues. A withdrawal period has not been established in pre-ruminating calves. Do not use in calves to be processed for veal.

**PRECAUTIONS:** The effects of Zuprevo 18% on bovine reproductive performance, pregnancy and lactation have not been determined. Swelling and inflammation, which may be severe, may be seen at the injection site after administration. Subcutaneous injection may result in local tissue reactions which persist beyond the slaughter withdrawal period. This may result in trim loss of edible tissue at slaughter.

Made in Germany  
Distributed by: Intervet Inc d/b/a  
Merck Animal Health, Summit,  
NJ 07901 Copyright © 2011, Intervet  
Inc., a subsidiary  
of Merck & Co. All rights reserved.

# NOT ALL VICTORY LAPS HAPPEN ON THE TRACK

© Copyright American Association of Bovine Practitioners; open access distribution.



**ZUPREVO™**  
(tildipirosin) **CHOOSE  
CONFIDENCE**

#### IMPORTANT SAFETY INFORMATION

FOR USE IN ANIMALS ONLY. NOT FOR HUMAN USE. KEEP OUT OF REACH OF CHILDREN. TO AVOID ACCIDENTAL INJECTION, DO NOT USE IN AUTOMATICALLY POWERED SYRINGES WHICH HAVE NO ADDITIONAL PROTECTION SYSTEM.

IN CASE OF HUMAN INJECTION, SEEK MEDICAL ADVICE IMMEDIATELY AND SHOW THE PACKAGE INSERT OR LABEL TO THE PHYSICIAN.

DO NOT USE Zuprevo™ 18% IN SWINE. Fatal adverse events have been reported following the use of tildipirosin in swine.  
NOT FOR USE IN CHICKENS OR TURKEYS.



© Copyright American Association of Bovine Practitioners. All rights reserved. Distribution of this advertisement is restricted to members of the American Association of Bovine Practitioners.

**Choose the BRD treatment that gets him back to the pen sooner.**  
**Defeat BRD with Zuprevo™ (tildipirosin), the treatment experts count on for fast-acting, long-lasting results.**  
**Ask your Merck Animal Health representative about Zuprevo. Learn more at [merck-animal-health-usa.com](http://merck-animal-health-usa.com).**  
**Choose Confidence. Choose Zuprevo.**

Cattle intended for human consumption must not be slaughtered within 21 days of the last treatment. Do not use in female dairy cattle 20 months of age or older. A withdrawal period has not been established in pre-ruminating calves. Do not use in calves to be processed for veal.

The effects Zuprevo™ 18% on bovine reproductive performance, pregnancy and lactation have not been determined. Swelling and inflammation, which may be severe, may be seen at the injection site after administration. Subcutaneous injection may result in local tissue reactions which persist beyond slaughter withdrawal period. This may result in trim loss of edible tissue at slaughter.

