

The Bovine Practitioner

Volume 56
Number 1
2022



The Official Publication
of the American Association
of Bovine Practitioners

BOVINE PRACTITIONER

Vol 56. NO 1.



Copy Editor

Geni Wren

1130 East Main St., Ste. 302
Ashland, OH 44805
800-269-2227

Advertising Representative

T. O'Brien & Assoc.

12118 Nieman Road
Overland Park, KS 66213
913-579-4084

Executive Editor

K. Fred Gingrich II, DVM

1130 East Main St., Ste. 302
Ashland, OH 44805
419-496-0685

Publisher

Truax Printing, Inc.

425 East Haskell St.
Loudonville, OH 44842
419-994-4166

Associate Editors

Dr. Virginia Fajt, dr.fajt@aabp.org

Dr. Miles Theurer, dr.theurer@aabp.org

Dr. Aurora Villarroel, dr.villarroel@aabp.org

Dr. Sarah Capik, dr.capik@aabp.org

The Bovine Practitioner is the official publication of The American Association of Bovine Practitioners. All manuscripts are peer-reviewed. *The Bovine Practitioner* is an open access journal.

©American Association of Bovine Practitioners, 2022 • All Rights Reserved
The Bovine Practitioner • ISSN 0524-1685

Table of Contents

Association of feedlot disease treatments on the probability of heart disease syndrome in U.S. feedlot cattle.....	1
<i>Blaine T. Johnson, David E. Amrine, Robert L. Larson, Brad J. White</i>	
Livestock fatalities attributed to a massive attack of dark rice field mosquitoes (<i>Psorophora columbiae</i>) following Hurricane	13
<i>Laura Bianca S. Cecco, Emi Sasaki, Daniel B. Paulsen, Craig Fontenot, Christine Navarre, Kristen Healy, Ingeborg M. Langohr</i>	
A mixed-method pilot study investigating challenges experienced by mixed animal veterinarians in practice and their private practice experiences prior to graduation	18
<i>Philippa M. Gibbons, Julia K. Wells, Kylie R.L. Watson, John J. Weale, Jerry R. Roberson</i>	
Nutritional management of calves grazing wheat and small grain pasture	29
<i>Paul A. Beck, R. Ryan Reuter</i>	
Associations between pen management characteristics and bovine respiratory disease incidence in the first 45 days post-arrival in feedlot cattle	40
<i>Hector A. Rojas, Brad J. White, David E. Amrine, Robert L. Larson, Sarah F. Capik</i>	
An evaluation of arrival metaphylaxis with enrofloxacin compared to tulathromycin in feedlot cattle at high risk of developing bovine respiratory disease in Mexico	53
<i>Luis O. Burciaga-Robles, Christopher McMullen, Ana Bras, Mariana Guerra-Maupome, Rodolfo A. Nava-Gaspar, Ryan D. Rademacher, Sherry J. Hannon, Victor E. Mercado-Talamantes, Calvin W. Booker</i>	
Comparison of the effect of different management strategies on the quality of recycled sand used for bedding in free stalls in southeastern Pennsylvania	60
<i>Emily E.A. Egolf, Michaela A. Kristula, Darko Stefanovski</i>	

Advertisers Index

Addison Biological Laboratory, Inc.	Inside front cover
Multimin USA, Inc.	Back cover

Notice to Readers

All statements, opinions and conclusions contained in articles in *The Bovine Practitioner* are those of the author(s), and are not necessarily those of the American Association of Bovine Practitioners (AABP) unless specifically approved by the AABP Board of Directors.

Pinkeye Problem? Pinkeye Solution!

MAXI/GUARD®

delivers twice the pinkeye
protection with
single dose convenience



Broad coverage

8 different *M. bovis* isolates
Over 90% efficacy

Easy syringeability

Approved for subcutaneous administration

Minimal injection site reactions

Strong safety record

Extremely low endotoxin levels

Single dose protection



Addison Biological Laboratory, Inc.

800-331-2530

www.addisonlabs.com

[f /AddisonLabs](https://www.facebook.com/AddisonLabs)





YOUR BEST SHOT AT IMMUNITY



Improve immune response to vaccines by 27%*

Multimin® 90 is a one-of-a-kind trace mineral injection that supports immune function. Research shows Multimin 90 in a calf vaccination program improves vaccine response from 53% to 80% as measured by a four-fold increase in antibody titers. Superior immune response provides better protection against BRD, while improving overall herd health and productivity. Optimize your vaccination ROI with Multimin 90. See corresponding page for prescribing information.

“MULTIMIN 90 is on top of our list... it has probably had the most dramatic effect on what we do in those protocols as anything we have done over the years.”

Dr. L.D. Barker and sons Dr. Matt Barker and Dr. Mark Barker
Professional Animal Health Center | Newcastle, OK

www.multiminusa.com

*Data on file.
© 2022 MULTIMIN® USA



KEEP OUT OF REACH OF CHILDREN

MULTIMIN® 90

An injectable aqueous supplemental source of zinc, manganese, selenium, and copper

CAUTION: FEDERAL LAW RESTRICTS THIS DRUG TO USE BY OR ON THE ORDER OF A LICENSED VETERINARIAN.

ACTIVE SUBSTANCES PER ML:

Zinc.....	60 mg/mL
Manganese.....	10 mg/mL
Selenium.....	5 mg/mL
Copper.....	15 mg/mL

OTHER SUBSTANCES:

Chlorocresol 0.1% w/v (as preservative)

DOSAGE RECOMMENDATIONS:

CALVES: Up to 1 year	1 mL/per 100 lbs. bodyweight
CATTLE: From 1-2 years	1 mL/per 150 lbs. bodyweight
CATTLE: Over 2 years	1 mL/per 200 lbs. bodyweight

PRECAUTION:

Selenium and copper are toxic if administered in excess.

Always follow recommended label dose. Do not overdose.

It is recommended that accurate body weight is determined prior to treatment.

Do not use concurrently with other injectable selenium and copper products.

Do not use concurrently with selenium or copper boluses.

Do not use in emaciated cattle with a BCS of 1 in dairy or 1-3 in beef.

Consult your veterinarian.

CAUTION:

Slight local reaction may occur for about 30 seconds after injection. A slight swelling may be observed at injection site for a few days after administration. Use standard aseptic procedures during administration of injections to reduce the risk of injection site abscesses or lesions.

WITHDRAWAL PERIOD:

Meat 14 days. Milk zero withdrawal.

DIRECTIONS:


This product is only for use in cattle.

MULTIMIN® 90 is to be given subcutaneously (under the skin) ONLY.

It is recommended to administer the product in accordance with Beef Quality Assurance (BQA) guidelines. Minimum distance between injection sites for the MULTIMIN® 90 product and other injection sites should be at least 4 inches.

Inject under the loose skin of the middle of the side of the neck. Max volume per injection site is 7 ml.

Subcutaneous injection in middle of side of neck.



Store Between 15°C and 30°C (59°F and 86°F).

Packaged in 100 mL & 500 mL size

NDC No. 49920-006-01
NDC No. 49920-006-05

SUPPLEMENTATION PROGRAM	
BULLS	3 times per year
BEEF COWS	4 weeks before breeding 4 weeks before calving
DAIRY COWS	4 weeks before calving 4 weeks before insemination at dry-off
CALVES	at birth at 3 months and/or weaning
HEIFERS	every 3 months - especially 4 weeks before breeding

(program gives planned dates that can be varied to suit management programs)

DOSAGE TABLE			
ANIMAL WEIGHT (lbs)	CALVES UP TO 1 YEAR 1 ml/100 lb BW	CATTLE 1 - 2 YEARS 1 ml/150 lb BW	CATTLE > 2 YEARS 1 ml/200 lb BW
50	0.5 ml	-	-
100	1 ml	-	-
150	1.5 ml	-	-
200	2 ml	-	-
300	3 ml	-	-
400	4 ml	-	-
500	5 ml	-	-
600	6 ml	-	-
700	7 ml	-	-
800	-	5.3 ml	-
900	-	6 ml	-
1000	-	6.6 ml	5 ml
1100	-	-	5.5 ml
1200	-	-	6 ml
1300	-	-	6.5 ml
1400	-	-	7 ml



MANUFACTURED FOR:
MULTIMIN®
NORTH AMERICA, INC.
Fort Collins, CO 80528